

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : C12N 15/82, 15/29, 5/10, C07K 14/415, C12Q 1/68		A2	(11) International Publication Number: WO 99/47689 (43) International Publication Date: 23 September 1999 (23.09.99)
(21) International Application Number: PCT/US99/05227 (22) International Filing Date: 10 March 1999 (10.03.99) (30) Priority Data: 60/078,144 16 March 1998 (16.03.98) US (71) Applicant (for all designated States except US): E.I. DU PONT DE NEMOURS AND COMPANY [US/US]; 1007 Market Street, Wilmington, DE 19898 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): CAHOON, Rebecca, E. [US/US]; 2331 West 18th Street, Wilmington, DE 19806 (US). CAIMI, Perry, G. [US/US]; 7 Holly Drive, Kennett Square, PA 19348 (US). KLEIN, Theodore, M. [US/US]; 2229 Rosewood Drive, Wilmington, DE 19810 (US). MORGANTE, Michele [IT/US]; 2306 Lorelei Lane, Wilmington, DE 19810 (US). SAKAI, Hajime [DE/US]; 105 Banbury Drive, Wilmington, DE 19803 (US). SHEN, Jennie, Bih-Jien [US/US]; 15 Bromley Court, Wilmington, DE 19810 (US). TINGEY, Scott, V. [US/US]; 35 Somerset Court, Elkton, MD 21921 (US).			(74) Agent: MAJARIAN, William, R.; E.I. du Pont de Nemours and Company, Legal Patent Records Center, 1007 Market Street, Wilmington, DE 19898 (US). (81) Designated States: AL, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GD, GE, HR, HU, ID, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: INHIBITORS OF APOPTOSIS PROTEINS IN PLANTS			
(57) Abstract This invention relates to an isolated nucleic acid fragment encoding an apoptosis inhibitory protein. The invention also relates to the construction of a chimeric gene encoding all or a portion of the apoptosis inhibitory protein, in sense or antisense orientation, wherein expression of the chimeric gene results in production of altered levels of the apoptosis inhibitory protein in a transformed host cell.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

TITLE

INHIBITORS OF APOPTOSIS PROTEINS IN PLANTS

FIELD OF THE INVENTION

5 This invention is in the field of plant molecular biology. More specifically, this invention pertains to nucleic acid fragments encoding proteins that inhibit apoptosis in plants and seeds.

BACKGROUND OF THE INVENTION

10 Multicellular organisms have evolved elaborate signal transduction pathways for maintaining homeostasis through the control of cell proliferation and death. Programmed cell death or apoptosis is a process in which unwanted cells are eliminated during growth and development (Steller, H. (1998) *PNAS* 95:5421-5422). Inhibitors of apoptosis proteins (IAPs) constitute a family of highly conserved death-suppressing proteins that were first
15 identified in baculoviruses (Liston, P. et al. (1997) *Apoptosis* 2(5):423-441). Two homologues have been identified in *Drosophila melanogaster* (D-IAP and D-IAP-2) and several in rodents and humans including human AAC-11 protein, X-linked inhibitor of apoptosis, XIAP and an XIAP associated protein ZAP-1 (Liston, P. et al. (1997) *Apoptosis* 2(5):423-441 and Liston, P. et al. (1996) *Nature* 379:349-352). Several DAD1 (defender
20 against death) proteins have also been identified in mammalian and plant cells (Gallosis, P. et al. (1997) *Plant J.* 11(6):1325-1331). These proteins appear to be a subunit of oligosaccharyltransferase (OST), an enzyme that is involved in N-linked glycosylation in eukaryotes and may also be associated with programmed cell death in mammalian and plant cells. No genes encoding IAP, IAP-2, ZAP-1 or AAC-11 proteins in plants, especially corn,
25 rice soybean and wheat, have been isolated and sequenced.

There is a great deal of interest in identifying the genes that encode IAPs in plants. These genes may be used to express IAPs in plant cells to enhance cell tissue culture growth. Accordingly, the availability of nucleic acid sequences encoding all or a portion of IAPs would facilitate studies to better understand programmed cell death in plants, provide genetic
30 tools to enhance cell growth in tissue culture, increase the efficiency of gene transfer and help provide more stable transformations.

SUMMARY OF THE INVENTION

The instant invention relates to isolated nucleic acid fragments encoding proteins that inhibit apoptosis. Specifically, this invention concerns an isolated nucleic acid fragment
35 encoding a DAD1 or AAC-11 protein. In addition, this invention relates to a nucleic acid fragment that is complementary to the nucleic acid fragment encoding a DAD1 or AAC-11 protein.

An additional embodiment of the instant invention pertains to a polypeptide encoding all or a substantial portion of an apoptosis inhibitory protein selected from the group consisting of DAD1 and AAC-11.

5 In another embodiment, the instant invention relates to a chimeric gene encoding a DAD1 or AAC-11 protein, or to a chimeric gene that comprises a nucleic acid fragment that is complementary to a nucleic acid fragment encoding a DAD1 or AAC-11 protein, operably linked to suitable regulatory sequences, wherein expression of the chimeric gene results in production of levels of the encoded protein in a transformed host cell that is altered (i.e., increased or decreased) from the level produced in an untransformed host cell.

10 In a further embodiment, the instant invention concerns a transformed host cell comprising in its genome a chimeric gene encoding a DAD1 or AAC-11 protein, operably linked to suitable regulatory sequences. Expression of the chimeric gene results in production of altered levels of the encoded protein in the transformed host cell. The transformed host cell can be of eukaryotic or prokaryotic origin, and include cells derived
15 from higher plants and microorganisms. The invention also includes transformed plants that arise from transformed host cells of higher plants, and seeds derived from such transformed plants.

An additional embodiment of the instant invention concerns a method of altering the level of expression of a DAD1 or AAC-11 protein in a transformed host cell comprising:
20 a) transforming a host cell with a chimeric gene comprising a nucleic acid fragment encoding a DAD1 or AAC-11 protein; and b) growing the transformed host cell under conditions that are suitable for expression of the chimeric gene wherein expression of the chimeric gene results in production of altered levels of DAD1 or AAC-11 protein in the transformed host cell.

25 An addition embodiment of the instant invention concerns a method for obtaining a nucleic acid fragment encoding all or a substantial portion of an amino acid sequence encoding a DAD1 or AAC-11 protein.

BRIEF DESCRIPTION OF THE FIGURES AND SEQUENCE LISTINGS

30 The invention can be more fully understood from the following detailed description and the accompanying Sequence Listing which form a part of this application.

Figure 1 shows a comparison of the amino acid sequences of the *Arabidopsis*, rice and pea DAD1 proteins (SEQ ID NO:13, SEQ ID NO:14 and SEQ ID NO:15, respectively) and
SEQ ID NOs:2, 4 and 6.

35 Figure 2 shows a comparison of the amino acid sequences of the human AAC-11 protein (SEQ ID NO:16) and SEQ ID NO:8.

The following sequence descriptions and Sequence Listing attached hereto comply with the rules governing nucleotide and/or amino acid sequence disclosures in patent applications as set forth in 37 C.F.R. §1.821-1.825.

SEQ ID NO:1 is the nucleotide sequence comprising the entire cDNA insert in clone plht.pk0012.c8 encoding a *Phaseolus* DAD1 protein.

SEQ ID NO:2 is the deduced amino acid sequence of an entire DAD1 protein derived from the nucleotide sequence of SEQ ID NO:1.

5 SEQ ID NO:3 is the nucleotide sequence comprising the entire cDNA insert in clone cbn10.pk0058.g12 encoding a corn DAD1 protein.

SEQ ID NO:4 is the deduced amino acid sequence of an entire DAD1 protein derived from the nucleotide sequence of SEQ ID NO:3.

10 SEQ ID NO:5 is the nucleotide sequence comprising a contig assembled from the cDNA inserts in clones sls1c.pk010.h9, ses2w.pk0011.h3, sdp4c.pk007.e17, se1.pk0035.e10 and sl1.pk0106.d1 encoding a soybean DAD1 protein.

SEQ ID NO:6 is the deduced amino acid sequence of an entire DAD1 protein derived from the nucleotide sequence of SEQ ID NO:5.

15 SEQ ID NO:7 is the nucleotide sequence comprising a contig assembled from the cDNA inserts in clones p0031.ccmba65r, p0037.crwaj44r, p0049.curar45r, p0004.cb1ha38r, p0011.cdapf47r, cbn10.pk0041.h3, cen1.pk0032.d4 and p0116.cesag38r encoding a corn AAC-11 protein.

SEQ ID NO:8 is the deduced amino acid sequence of an entire AAC-11 protein derived from the nucleotide sequence of SEQ ID NO:7.

20 SEQ ID NO:9 is the nucleotide sequence comprising a portion of the cDNA insert in clone rls24.pk0025.b4 encoding a rice AAC-11 protein.

SEQ ID NO:10 is the deduced amino acid sequence of a portion of an AAC-11 protein derived from the nucleotide sequence of SEQ ID NO:9.

25 SEQ ID NO:11 is the nucleotide sequence comprising a contig assembled from the cDNA inserts in clones sdp2c.pk027.j3 and src3c.pk001.j10 encoding a soybean AAC-11 protein.

SEQ ID NO:12 is the deduced amino acid sequence of a portion of a AAC-11 protein derived from the nucleotide sequence of SEQ ID NO:11.

30 SEQ ID NO:13 is the amino acid sequence of the *Arabidopsis thaliana* DAD1 protein set forth in NCBI Identifier No. gi 2623638.

SEQ ID NO:14 is the amino acid sequence of the *Oryza sativa* DAD1 protein set forth in NCBI Identifier No. gi 2723473.

SEQ ID NO:15 is the amino acid sequence of the *Pisum sativum* DAD1 protein set forth in NCBI Identifier No. gi 3869255.

35 SEQ ID NO:16 is the amino acid sequence of the *Homo sapiens* AAC-11 protein set forth in NCBI Identifier No. gi 2623761.

The Sequence Listing contains the one letter code for nucleotide sequence characters and the three letter codes for amino acids as defined in conformity with the IUPAC-IUBMB standards described in *Nucleic Acids Research* 13:3021-3030 (1985) and in the *Biochemical*

Journal 219 (No. 2):345-373 (1984) which are herein incorporated by reference. The symbols and format used for nucleotide and amino acid sequence data comply with the rules set forth in 37 C.F.R. §1.822.

DETAILED DESCRIPTION OF THE INVENTION

5 In the context of this disclosure, a number of terms shall be utilized. As used herein, an "isolated nucleic acid fragment" is a polymer of RNA or DNA that is single- or double-stranded, optionally containing synthetic, non-natural or altered nucleotide bases. An isolated nucleic acid fragment in the form of a polymer of DNA may be comprised of one or more segments of cDNA, genomic DNA or synthetic DNA. As used herein, "contig" refers to an assemblage of overlapping nucleic acid sequences to form one contiguous nucleotide sequence. For example, several DNA sequences can be compared and aligned to identify common or overlapping regions. The individual sequences can then be assembled into a single contiguous nucleotide sequence. As used herein a "peptide", "polypeptide" or protein refers to a linear polymer composed of amino acids connected by peptide bonds.

15 As used herein, "substantially similar" refers to nucleic acid fragments wherein changes in one or more nucleotide bases results in substitution of one or more amino acids, but do not affect the functional properties of the protein encoded by the DNA sequence. "Substantially similar" also refers to nucleic acid fragments wherein changes in one or more nucleotide bases does not affect the ability of the nucleic acid fragment to mediate alteration of gene expression by antisense or co-suppression technology. "Substantially similar" also refers to modifications of the nucleic acid fragments of the instant invention such as deletion or insertion of one or more nucleotides that do not substantially affect the functional properties of the resulting transcript vis-à-vis the ability to mediate alteration of gene expression by antisense or co-suppression technology or alteration of the functional properties of the resulting protein molecule. It is therefore understood that the invention encompasses more than the specific exemplary sequences.

25 For example, it is well known in the art that antisense suppression and co-suppression of gene expression may be accomplished using nucleic acid fragments representing less than the entire coding region of a gene, and by nucleic acid fragments that do not share 100% sequence identity with the gene to be suppressed. Moreover, alterations in a gene which result in the production of a chemically equivalent amino acid at a given site, but do not effect the functional properties of the encoded protein, are well known in the art. Thus, a codon for the amino acid alanine, a hydrophobic amino acid, may be substituted by a codon encoding another less hydrophobic residue, such as glycine, or a more hydrophobic residue, such as valine, leucine, or isoleucine. Similarly, changes which result in substitution of one negatively charged residue for another, such as aspartic acid for glutamic acid, or one positively charged residue for another, such as lysine for arginine, can also be expected to produce a functionally equivalent product. Nucleotide changes which result in alteration of the N-terminal and C-terminal portions of the protein molecule would also not be expected

to alter the activity of the protein. Each of the proposed modifications is well within the routine skill in the art, as is determination of retention of biological activity of the encoded products.

Moreover, substantially similar nucleic acid fragments may also be characterized by their ability to hybridize, under stringent conditions (0.1X SSC, 0.1% SDS, 65°C), with the nucleic acid fragments disclosed herein.

Substantially similar nucleic acid fragments of the instant invention may also be characterized by the percent similarity of the amino acid sequences that they encode to the amino acid sequences disclosed herein, as determined by algorithms commonly employed by those skilled in this art. Preferred are those nucleic acid fragments whose nucleotide sequences encode amino acid sequences that are 85% similar to the amino acid sequences reported herein. More preferred nucleic acid fragments encode amino acid sequences that are 90% similar to the amino acid sequences reported herein. Most preferred are nucleic acid fragments that encode amino acid sequences that are 95% similar to the amino acid sequences reported herein. Sequence alignments and percent similarity calculations were performed using the Megalign program of the LASARGENE bioinformatics computing suite (DNASTAR Inc., Madison, WI). Multiple alignment of the sequences was performed using the Clustal method of alignment (Higgins, D.G. and Sharp, P.M. (1989) *CABIOS* 5:151-153) with the default parameters (GAP PENALTY=10, GAP LENGTH PENALTY=10). Default parameters for pairwise alignments using the Clustal method were KTUPLE 1, GAP PENALTY=3, WINDOW=5 and DIAGONALS SAVED=5.

A "substantial portion" of an amino acid or nucleotide sequence comprises enough of the amino acid sequence of a polypeptide or the nucleotide sequence of a gene to afford putative identification of that polypeptide or gene, either by manual evaluation of the sequence by one skilled in the art, or by computer-automated sequence comparison and identification using algorithms such as BLAST (Basic Local Alignment Search Tool; Altschul, S. F., et al., (1993) *J. Mol. Biol.* 215:403-410; see also www.ncbi.nlm.nih.gov/BLAST/). In general, a sequence of ten or more contiguous amino acids or thirty or more nucleotides is necessary in order to putatively identify a polypeptide or nucleic acid sequence as homologous to a known protein or gene. Moreover, with respect to nucleotide sequences, gene specific oligonucleotide probes comprising 20-30 contiguous nucleotides may be used in sequence-dependent methods of gene identification (e.g., Southern hybridization) and isolation (e.g., *in situ* hybridization of bacterial colonies or bacteriophage plaques). In addition, short oligonucleotides of 12-15 bases may be used as amplification primers in PCR in order to obtain a particular nucleic acid fragment comprising the primers. Accordingly, a "substantial portion" of a nucleotide sequence comprises enough of the sequence to afford specific identification and/or isolation of a nucleic acid fragment comprising the sequence. The instant specification teaches partial or complete amino acid and nucleotide sequences encoding one or more particular plant

proteins. The skilled artisan, having the benefit of the sequences as reported herein, may now use all or a substantial portion of the disclosed sequences for purposes known to those skilled in this art. Accordingly, the instant invention comprises the complete sequences as reported in the accompanying Sequence Listing, as well as substantial portions of those sequences as defined above.

"Codon degeneracy" refers to divergence in the genetic code permitting variation of the nucleotide sequence without effecting the amino acid sequence of an encoded polypeptide. Accordingly, the instant invention relates to any nucleic acid fragment that encodes all or a substantial portion of the amino acid sequence encoding the DAD1 or AAC-11 proteins as set forth in SEQ ID NOs:2, 4, 6, 8, 10, 12, 14, 16 and 18. The skilled artisan is well aware of the "codon-bias" exhibited by a specific host cell in usage of nucleotide codons to specify a given amino acid. Therefore, when synthesizing a gene for improved expression in a host cell, it is desirable to design the gene such that its frequency of codon usage approaches the frequency of preferred codon usage of the host cell.

"Synthetic genes" can be assembled from oligonucleotide building blocks that are chemically synthesized using procedures known to those skilled in the art. These building blocks are ligated and annealed to form gene segments which are then enzymatically assembled to construct the entire gene. "Chemically synthesized", as related to a sequence of DNA, means that the component nucleotides were assembled *in vitro*. Manual chemical synthesis of DNA may be accomplished using well established procedures, or automated chemical synthesis can be performed using one of a number of commercially available machines. Accordingly, the genes can be tailored for optimal gene expression based on optimization of nucleotide sequence to reflect the codon bias of the host cell. The skilled artisan appreciates the likelihood of successful gene expression if codon usage is biased towards those codons favored by the host. Determination of preferred codons can be based on a survey of genes derived from the host cell where sequence information is available.

"Gene" refers to a nucleic acid fragment that expresses a specific protein, including regulatory sequences preceding (5' non-coding sequences) and following (3' non-coding sequences) the coding sequence. "Native gene" refers to a gene as found in nature with its own regulatory sequences. "Chimeric gene" refers any gene that is not a native gene, comprising regulatory and coding sequences that are not found together in nature. Accordingly, a chimeric gene may comprise regulatory sequences and coding sequences that are derived from different sources, or regulatory sequences and coding sequences derived from the same source, but arranged in a manner different than that found in nature.

"Endogenous gene" refers to a native gene in its natural location in the genome of an organism. A "foreign" gene refers to a gene not normally found in the host organism, but that is introduced into the host organism by gene transfer. Foreign genes can comprise native genes inserted into a non-native organism, or chimeric genes. A "transgene" is a gene that has been introduced into the genome by a transformation procedure.

“Coding sequence” refers to a DNA sequence that codes for a specific amino acid sequence. “Regulatory sequences” refer to nucleotide sequences located upstream (5' non-coding sequences), within, or downstream (3' non-coding sequences) of a coding sequence, and which influence the transcription, RNA processing or stability, or translation of the associated coding sequence. Regulatory sequences may include promoters, translation leader sequences, introns, and polyadenylation recognition sequences.

“Promoter” refers to a DNA sequence capable of controlling the expression of a coding sequence or functional RNA. In general, a coding sequence is located 3' to a promoter sequence. The promoter sequence consists of proximal and more distal upstream elements, the latter elements often referred to as enhancers. Accordingly, an “enhancer” is a DNA sequence which can stimulate promoter activity and may be an innate element of the promoter or a heterologous element inserted to enhance the level or tissue-specificity of a promoter. Promoters may be derived in their entirety from a native gene, or be composed of different elements derived from different promoters found in nature, or even comprise synthetic DNA segments. It is understood by those skilled in the art that different promoters may direct the expression of a gene in different tissues or cell types, or at different stages of development, or in response to different environmental conditions. Promoters which cause a gene to be expressed in most cell types at most times are commonly referred to as “constitutive promoters”. New promoters of various types useful in plant cells are constantly being discovered; numerous examples may be found in the compilation by Okamuro and Goldberg, (1989) *Biochemistry of Plants* 15:1-82. It is further recognized that since in most cases the exact boundaries of regulatory sequences have not been completely defined, DNA fragments of different lengths may have identical promoter activity.

The “translation leader sequence” refers to a DNA sequence located between the promoter sequence of a gene and the coding sequence. The translation leader sequence is present in the fully processed mRNA upstream of the translation start sequence. The translation leader sequence may affect processing of the primary transcript to mRNA, mRNA stability or translation efficiency. Examples of translation leader sequences have been described (Turner, R. and Foster, G.D. (1995) *Molecular Biotechnology* 3:225).

The “3' non-coding sequences” refer to DNA sequences located downstream of a coding sequence and include polyadenylation recognition sequences and other sequences encoding regulatory signals capable of affecting mRNA processing or gene expression. The polyadenylation signal is usually characterized by affecting the addition of polyadenylic acid tracts to the 3' end of the mRNA precursor. The use of different 3' non-coding sequences is exemplified by Ingelbrecht et al., (1989) *Plant Cell* 1:671-680.

“RNA transcript” refers to the product resulting from RNA polymerase-catalyzed transcription of a DNA sequence. When the RNA transcript is a perfect complementary copy of the DNA sequence, it is referred to as the primary transcript or it may be a RNA sequence derived from posttranscriptional processing of the primary transcript and is

referred to as the mature RNA. "Messenger RNA (mRNA)" refers to the RNA that is without introns and that can be translated into protein by the cell. "cDNA" refers to a double-stranded DNA that is complementary to and derived from mRNA. "Sense" RNA refers to RNA transcript that includes the mRNA and so can be translated into protein by the cell. "Antisense RNA" refers to a RNA transcript that is complementary to all or part of a target primary transcript or mRNA and that blocks the expression of a target gene (U.S. Pat. No. 5,107,065, incorporated herein by reference). The complementarity of an antisense RNA may be with any part of the specific gene transcript, i.e., at the 5' non-coding sequence, 3' non-coding sequence, introns, or the coding sequence. "Functional RNA" refers to sense RNA, antisense RNA, ribozyme RNA, or other RNA that may not be translated but yet has an effect on cellular processes.

The term "operably linked" refers to the association of nucleic acid sequences on a single nucleic acid fragment so that the function of one is affected by the other. For example, a promoter is operably linked with a coding sequence when it is capable of affecting the expression of that coding sequence (i.e., that the coding sequence is under the transcriptional control of the promoter). Coding sequences can be operably linked to regulatory sequences in sense or antisense orientation.

The term "expression", as used herein, refers to the transcription and stable accumulation of sense (mRNA) or antisense RNA derived from the nucleic acid fragment of the invention. Expression may also refer to translation of mRNA into a polypeptide. "Antisense inhibition" refers to the production of antisense RNA transcripts capable of suppressing the expression of the target protein. "Overexpression" refers to the production of a gene product in transgenic organisms that exceeds levels of production in normal or non-transformed organisms. "Co-suppression" refers to the production of sense RNA transcripts capable of suppressing the expression of identical or substantially similar foreign or endogenous genes (U.S. Pat. No. 5,231,020, incorporated herein by reference).

"Altered levels" refers to the production of gene product(s) in transgenic organisms in amounts or proportions that differ from that of normal or non-transformed organisms.

"Mature" protein refers to a post-translationally processed polypeptide; i.e., one from which any pre- or propeptides present in the primary translation product have been removed. "Precursor" protein refers to the primary product of translation of mRNA; i.e., with pre- and propeptides still present. Pre- and propeptides may be but are not limited to intracellular localization signals.

A "chloroplast transit peptide" is an amino acid sequence which is translated in conjunction with a protein and directs the protein to the chloroplast or other plastid types present in the cell in which the protein is made. "Chloroplast transit sequence" refers to a nucleotide sequence that encodes a chloroplast transit peptide. A "signal peptide" is an amino acid sequence which is translated in conjunction with a protein and directs the protein to the secretory system (Chrispeels, J. J., (1991) *Ann. Rev. Plant Phys. Plant Mol. Biol.*

42:21-53). If the protein is to be directed to a vacuole, a vacuolar targeting signal (*supra*) can further be added, or if to the endoplasmic reticulum, an endoplasmic reticulum retention signal (*supra*) may be added. If the protein is to be directed to the nucleus, any signal peptide present should be removed and instead a nuclear localization signal included (Raikhel (1992) *Plant Phys.* 100:1627-1632).

"Transformation" refers to the transfer of a nucleic acid fragment into the genome of a host organism, resulting in genetically stable inheritance. Host organisms containing the transformed nucleic acid fragments are referred to as "transgenic" organisms. Examples of methods of plant transformation include *Agrobacterium*-mediated transformation (De Blaere et al. (1987) *Meth. Enzymol.* 143:277) and particle-accelerated or "gene gun" transformation technology (Klein et al. (1987) *Nature (London)* 327:70-73; U.S. Pat. No. 4,945,050, incorporated herein by reference).

Standard recombinant DNA and molecular cloning techniques used herein are well known in the art and are described more fully in Sambrook, J., Fritsch, E.F. and Maniatis, T. *Molecular Cloning: A Laboratory Manual*; Cold Spring Harbor Laboratory Press: Cold Spring Harbor, 1989 (hereinafter "Maniatis").

Nucleic acid fragments encoding at least a portion of several apoptosis inhibitory proteins have been isolated and identified by comparison of random plant cDNA sequences to public databases containing nucleotide and protein sequences using the BLAST algorithms well known to those skilled in the art. Table 1 lists the proteins that are described herein, and the designation of the cDNA clones that comprise the nucleic acid fragments encoding these proteins.

TABLE 1
Apoptosis Inhibitory Proteins

Enzyme	Clone	Plant
Defenders against cell death protein DAD1	plht.pk0012.c8	Lima bean
	cbn10.pk0058.g12	Maize
	sls1c.pk010.h9	Soybean
	ses2w.pk0011.h3	Soybean
	sdp4c.pk007.e17	Soybean
	se1.pk0035.e10	Soybean
	sl1.pk0106.d1	Soybean

Enzyme	Clone	Plant
Inhibitor of Apoptosis Protein AAC-11	p0031.ccmba65r	Corn
	p0037.crwaj44r	Corn
	p0049.curar45r	Corn
	p0004.cb1ha38r	Corn
	p0011.cdapf47r	Corn
	cbn10.pk0041.h3	Corn
	cen1.pk0032.d4	Corn
	p0116.cesag38r	Corn
	rls24.pk0025.b4	Rice
	sdp2c.pk027.j3	Soybean
	src3c.pk001.j10	Soybean

The nucleic acid fragments of the instant invention may be used to isolate cDNAs and genes encoding homologous proteins from the same or other plant species. Isolation of homologous genes using sequence-dependent protocols is well known in the art. Examples of sequence-dependent protocols include, but are not limited to, methods of nucleic acid hybridization, and methods of DNA and RNA amplification as exemplified by various uses of nucleic acid amplification technologies (e.g., polymerase chain reaction, ligase chain reaction).

For example, genes encoding other DAD1 or AAC-11 protein, either as cDNAs or genomic DNAs, could be isolated directly by using all or a portion of the instant nucleic acid fragments as DNA hybridization probes to screen libraries from any desired plant employing methodology well known to those skilled in the art. Specific oligonucleotide probes based upon the instant nucleic acid sequences can be designed and synthesized by methods known in the art (Maniatis). Moreover, the entire sequences can be used directly to synthesize DNA probes by methods known to the skilled artisan such as random primer DNA labeling, nick translation, or end-labeling techniques, or RNA probes using available *in vitro* transcription systems. In addition, specific primers can be designed and used to amplify a part or all of the instant sequences. The resulting amplification products can be labeled directly during amplification reactions or labeled after amplification reactions, and used as probes to isolate full length cDNA or genomic fragments under conditions of appropriate stringency.

In addition, two short segments of the instant nucleic acid fragments may be used in polymerase chain reaction protocols to amplify longer nucleic acid fragments encoding homologous genes from DNA or RNA. The polymerase chain reaction may also be performed on a library of cloned nucleic acid fragments wherein the sequence of one primer is derived from the instant nucleic acid fragments, and the sequence of the other primer takes advantage of the presence of the polyadenylic acid tracts to the 3' end of the mRNA

precursor encoding plant genes. Alternatively, the second primer sequence may be based upon sequences derived from the cloning vector. For example, the skilled artisan can follow the RACE protocol (Frohman et al., (1988) *PNAS USA* 85:8998) to generate cDNAs by using PCR to amplify copies of the region between a single point in the transcript and the 3' or 5' end. Primers oriented in the 3' and 5' directions can be designed from the instant sequences. Using commercially available 3' RACE or 5' RACE systems (BRL), specific 3' or 5' cDNA fragments can be isolated (Ohara et al., (1989) *PNAS USA* 86:5673; Loh et al., (1989) *Science* 243:217). Products generated by the 3' and 5' RACE procedures can be combined to generate full-length cDNAs (Frohman, M. A. and Martin, G. R., (1989)

10 *Techniques* 1:165).

Availability of the instant nucleotide and deduced amino acid sequences facilitates immunological screening of cDNA expression libraries. Synthetic peptides representing portions of the instant amino acid sequences may be synthesized. These peptides can be used to immunize animals to produce polyclonal or monoclonal antibodies with specificity for peptides or proteins comprising the amino acid sequences. These antibodies can be then be used to screen cDNA expression libraries to isolate full-length cDNA clones of interest (Lerner, R. A. (1984) *Adv. Immunol.* 36:1; Maniatis).

The nucleic acid fragments of the instant invention may be used to create transgenic plants in which the disclosed DAD1 or AAC-11 proteins are present at higher or lower levels than normal or in cell types or developmental stages in which they are not normally found. This would have the effect of altering the level of DAD1 or AAC-11 proteins in those cells. Altering the level of DAD1 and AAC-11 proteins in cells could facilitate studies to better understand programmed cell death in plants, provide genetic tools to alter programmed cell death, enhance cell growth in tissue culture, increase the efficiency of gene transfer and help provide more stable transformations.

Overexpression of the DAD1 or AAC-11 proteins of the instant invention may be accomplished by first constructing a chimeric gene in which the coding region is operably linked to a promoter capable of directing expression of a gene in the desired tissues at the desired stage of development. For reasons of convenience, the chimeric gene may comprise promoter sequences and translation leader sequences derived from the same genes. 3' Non-coding sequences encoding transcription termination signals may also be provided. The instant chimeric gene may also comprise one or more introns in order to facilitate gene expression.

Plasmid vectors comprising the instant chimeric gene can then be constructed. The choice of plasmid vector is dependent upon the method that will be used to transform host plants. The skilled artisan is well aware of the genetic elements that must be present on the plasmid vector in order to successfully transform, select and propagate host cells containing the chimeric gene. The skilled artisan will also recognize that different independent transformation events will result in different levels and patterns of expression (Jones et al.,

(1985) *EMBO J.* 4:2411-2418; De Almeida et al., (1989) *Mol. Gen. Genetics* 218:78-86), and thus that multiple events must be screened in order to obtain lines displaying the desired expression level and pattern. Such screening may be accomplished by Southern analysis of DNA, Northern analysis of mRNA expression, Western analysis of protein expression, or phenotypic analysis.

For some applications it may be useful to direct the instant apoptosis inhibitory proteins to different cellular compartments, or to facilitate its secretion from the cell. It is thus envisioned that the chimeric gene described above may be further supplemented by altering the coding sequence to encode a DAD1 or AAC-11 protein with appropriate intracellular targeting sequences such as transit sequences (Keegstra, K. (1989) *Cell* 56:247-253), signal sequences or sequences encoding endoplasmic reticulum localization (Chrispeels, J. J., (1991) *Ann. Rev. Plant Phys. Plant Mol. Biol.* 42:21-53), or nuclear localization signals (Raikhel, N. (1992) *Plant Phys.* 100:1627-1632) added and/or with targeting sequences that are already present removed. While the references cited give examples of each of these, the list is not exhaustive and more targeting signals of utility may be discovered in the future.

It may also be desirable to reduce or eliminate expression of genes encoding DAD1 or AAC-11 proteins in plants for some applications. In order to accomplish this, a chimeric gene designed for co-suppression of the instant apoptosis inhibitory proteins can be constructed by linking a gene or gene fragment encoding a DAD1 or AAC-11 protein to plant promoter sequences. Alternatively, a chimeric gene designed to express antisense RNA for all or part of the instant nucleic acid fragment can be constructed by linking the gene or gene fragment in reverse orientation to plant promoter sequences. Either the co-suppression or antisense chimeric genes could be introduced into plants via transformation wherein expression of the corresponding endogenous genes are reduced or eliminated.

The instant DAD1 or AAC-11 proteins (or portions thereof) may be produced in heterologous host cells, particularly in the cells of microbial hosts, and can be used to prepare antibodies to the these proteins by methods well known to those skilled in the art. The antibodies are useful for detecting DAD1 or AAC-11 proteins *in situ* in cells or *in vitro* in cell extracts. Preferred heterologous host cells for production of the instant DAD1 or AAC-11 proteins are microbial hosts. Microbial expression systems and expression vectors containing regulatory sequences that direct high level expression of foreign proteins are well known to those skilled in the art. Any of these could be used to construct a chimeric gene for production of the instant DAD1 or AAC-11 proteins. This chimeric gene could then be introduced into appropriate microorganisms via transformation to provide high level expression of the encoded apoptosis inhibitory protein. An example of a vector for high level expression of the instant DAD1 or AAC-11 proteins in a bacterial host is provided (Example 7).

All or a substantial portion of the nucleic acid fragments of the instant invention may also be used as probes for genetically and physically mapping the genes that they are a part of, and as markers for traits linked to those genes. Such information may be useful in plant breeding in order to develop lines with desired phenotypes. For example, the instant nucleic acid fragments may be used as restriction fragment length polymorphism (RFLP) markers. Southern blots (Maniatis) of restriction-digested plant genomic DNA may be probed with the nucleic acid fragments of the instant invention. The resulting banding patterns may then be subjected to genetic analyses using computer programs such as MapMaker (Lander et al., (1987) *Genomics* 1:174-181) in order to construct a genetic map. In addition, the nucleic acid fragments of the instant invention may be used to probe Southern blots containing restriction endonuclease-treated genomic DNAs of a set of individuals representing parent and progeny of a defined genetic cross. Segregation of the DNA polymorphisms is noted and used to calculate the position of the instant nucleic acid sequence in the genetic map previously obtained using this population (Botstein, D. et al., (1980) *Am. J. Hum. Genet.* 32:314-331).

The production and use of plant gene-derived probes for use in genetic mapping is described in R. Bernatzky, R. and Tanksley, S. D. (1986) *Plant Mol. Biol. Reporter* 4(1):37-41. Numerous publications describe genetic mapping of specific cDNA clones using the methodology outlined above or variations thereof. For example, F2 intercross populations, backcross populations, randomly mated populations, near isogenic lines, and other sets of individuals may be used for mapping. Such methodologies are well known to those skilled in the art.

Nucleic acid probes derived from the instant nucleic acid sequences may also be used for physical mapping (i.e., placement of sequences on physical maps; see Hoheisel, J. D., et al., In: *Nonmammalian Genomic Analysis: A Practical Guide*, Academic press 1996, pp. 319-346, and references cited therein).

In another embodiment, nucleic acid probes derived from the instant nucleic acid sequences may be used in direct fluorescence *in situ* hybridization (FISH) mapping (Trask, B. J. (1991) *Trends Genet.* 7:149-154). Although current methods of FISH mapping favor use of large clones (several to several hundred KB; see Laan, M. et al. (1995) *Genome Research* 5:13-20), improvements in sensitivity may allow performance of FISH mapping using shorter probes.

A variety of nucleic acid amplification-based methods of genetic and physical mapping may be carried out using the instant nucleic acid sequences. Examples include allele-specific amplification (Kazazian, H. H. (1989) *J. Lab. Clin. Med.* 114(2):95-96), polymorphism of PCR-amplified fragments (CAPS; Sheffield, V. C. et al. (1993) *Genomics* 16:325-332), allele-specific ligation (Landegren, U. et al. (1988) *Science* 241:1077-1080), nucleotide extension reactions (Sokolov, B. P. (1990) *Nucleic Acid Res.* 18:3671), Radiation Hybrid Mapping (Walter, M. A. et al. (1997) *Nature Genetics* 7:22-28) and Happy Mapping

(Dear, P. H. and Cook, P. R. (1989) *Nucleic Acid Res.* 17:6795-6807). For these methods, the sequence of a nucleic acid fragment is used to design and produce primer pairs for use in the amplification reaction or in primer extension reactions. The design of such primers is well known to those skilled in the art. In methods employing PCR-based genetic mapping, it may be necessary to identify DNA sequence differences between the parents of the mapping cross in the region corresponding to the instant nucleic acid sequence. This, however, is generally not necessary for mapping methods.

Loss of function mutant phenotypes may be identified for the instant cDNA clones either by targeted gene disruption protocols or by identifying specific mutants for these genes contained in a maize population carrying mutations in all possible genes (Ballinger and Benzer, (1989) *Proc. Natl. Acad. Sci USA* 86:9402; Koes et al., (1995) *Proc. Natl. Acad. Sci USA* 92:8149; Bensen et al., (1995) *Plant Cell* 7:75). The latter approach may be accomplished in two ways. First, short segments of the instant nucleic acid fragments may be used in polymerase chain reaction protocols in conjunction with a mutation tag sequence primer on DNAs prepared from a population of plants in which Mutator transposons or some other mutation-causing DNA element has been introduced (see Bensen, *supra*). The amplification of a specific DNA fragment with these primers indicates the insertion of the mutation tag element in or near the plant gene encoding the DAD1 or AAC-11 protein. Alternatively, the instant nucleic acid fragment may be used as a hybridization probe against PCR amplification products generated from the mutation population using the mutation tag sequence primer in conjunction with an arbitrary genomic site primer, such as that for a restriction enzyme site-anchored synthetic adaptor. With either method, a plant containing a mutation in the endogenous gene encoding a DAD1 or AAC-11 protein can be identified and obtained. This mutant plant can then be used to determine or confirm the natural function of the DAD1 or AAC-11 protein gene product.

EXAMPLES

The present invention is further defined in the following Examples, in which all parts and percentages are by weight and degrees are Celsius, unless otherwise stated. It should be understood that these Examples, while indicating preferred embodiments of the invention, are given by way of illustration only. From the above discussion and these Examples, one skilled in the art can ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions.

EXAMPLE 1

Composition of cDNA Libraries; Isolation and Sequencing of cDNA Clones

cDNA libraries representing mRNAs from various corn, rice, *Phaseolus* and soybean tissues were prepared. The characteristics of the libraries are described below.

TABLE 2
cDNA Libraries from Corn, Rice, *Phaseolus* and Soybean

Library	Tissue	Clone
cbn10	Corn developing kernel (embryo and endosperm; 10 days after pollination)	cbn10.pk0058.g12 cbn10.pk0041.h3
cen1	Corn endosperm 11 days after pollination	cen1.pk0032.d4
p0031	Corn shoot culture, initiated from seed derived meristems	p0031.ccmba65r
p0037	Corn, V5 roots, infested with corn root worm	p0037.crwaj44r
p0049	Corn, 5 days after pollination, whole kernels	p0049.curar45r
p0004	Corn immature ear	p0004.cb1ha38r
p0011	Corn, 5 days after pollination whole kernels	p0011.cdapf47r
p0116	Corn, DAM methylase induced transgenic suspension cells*	p0116.cesag38r
plht	<i>Phaseolus lunatus</i> leaf- heat tolerant	plht.pk0012.c8
rls24	Rice leaf (15 days after pollination) 24 hours after infection of <i>Magaporthe grisea</i> strain 4360-R-67 (avr2-yamo); Susceptible	rls24.pk0025.b4
sdp4c	Soybean (<i>Glycine max</i> L.) developing embryo (9-11 mm)	sdp4c.pk007.e17
sdp2c	Soybean (<i>Glycine max</i> L.) developing pods 6-7 mm	sdp2c.pk027.j3
ses2w	Soybean (<i>Glycine max</i> L.) embryogenic suspension 2 weeks after subculture	ses2w.pk0011.h3
se1	Soybean (<i>Glycine max</i> L.) embryo, 6-10 days after flowering	se1.pk0035.e10
sl1	Soybean (<i>Glycine max</i> L.) two week old developing seedlings treated with water	sl1.pk0106.d1
sls1c	Soybean (<i>Glycine max</i> L., S1990) infected with <i>Sclerotinia sclerotiorum</i> mycelium	sls1c.pk010.h9
src3c	Soybean (<i>Glycine max</i> L., Bell) 8 day old root inoculated with eggs of cyst nematode <i>Heterodera glycines</i> (Race 14) for 4 days	src3c.pk001.j10

- 5 *Cell line is transgenic for a vector harboring four copies of the estrogen response element (ERE) and CaMV -59 promoter driving dam methylase expression (Klein-Hitpab, L., et al., (1989) *Cell* 46:1053-1061). Expression of dam methylase was induced by 17 alpha-ethnylestradiol.

- 10 cDNA libraries were prepared in Uni-ZAP™ XR vectors according to the manufacturer's protocol (Stratagene Cloning Systems, La Jolla, CA). Conversion of the Uni-ZAP™ XR libraries into plasmid libraries was accomplished according to the protocol provided by Stratagene. Upon conversion, cDNA inserts were contained in the plasmid vector pBluescript. cDNA inserts from randomly picked bacterial colonies containing recombinant pBluescript plasmids were amplified via polymerase chain reaction using primers specific for vector sequences flanking the inserted cDNA sequences or plasmid
- 15 DNA was prepared from cultured bacterial cells. Amplified insert DNAs or plasmid DNAs were sequenced in dye-primer sequencing reactions to generate partial cDNA sequences

(expressed sequence tags or "ESTs"; see Adams, M. D. et al., (1991) *Science* 252:1651). The resulting ESTs were analyzed using a Perkin Elmer Model 377 fluorescent sequencer.

EXAMPLE 2

Identification of cDNA Clones

5 ESTs encoding apoptosis inhibitory proteins were identified by conducting BLAST (Basic Local Alignment Search Tool; Altschul, S. F., et al., (1993) *J. Mol. Biol.* 215:403-410; see also www.ncbi.nlm.nih.gov/BLAST/) searches for similarity to sequences contained in the BLAST "nr" database (comprising all non-redundant GenBank CDS translations, sequences derived from the 3-dimensional structure Brookhaven Protein

10 Data Bank, the last major release of the SWISS-PROT protein sequence database, EMBL, and DDBJ databases). The cDNA sequences obtained in Example 1 were analyzed for similarity to all publicly available DNA sequences contained in the "nr" database using the BLASTN algorithm provided by the National Center for Biotechnology Information (NCBI). The DNA sequences were translated in all reading frames and compared for

15 similarity to all publicly available protein sequences contained in the "nr" database using the BLASTX algorithm (Gish, W. and States, D. J. (1993) *Nature Genetics* 3:266-272 and Altschul, Stephen F., et al. (1997) *Nucleic Acids Res.* 25:3389-3402) provided by the NCBI. For convenience, the P-value (probability) of observing a match of a cDNA sequence to a sequence contained in the searched databases merely by chance as calculated by BLAST

20 are reported herein as "pLog" values, which represent the negative of the logarithm of the reported P-value. Accordingly, the greater the pLog value, the greater the likelihood that the cDNA sequence and the BLAST "hit" represent homologous proteins.

EXAMPLE 3

Characterization of cDNA Clones Encoding DAD1 Homologs

25 The BLASTX search using the EST sequences from clones plht.pk0012.c8 and sel.pk0035.e10 revealed similarity of the proteins encoded by the cDNAs to DAD1 from *Arabidopsis thaliana* (NCBI Identifier No. gi 3608136). Subsequently, several other soybean clones (sls1c.pk010.h9, ses2w.pk0011.h3, sdp4c.pk007.e17 and sl1.pk0106.d1) were obtained that had similarity of the proteins encoded by the cDNAs to DAD1 from

30 *Arabidopsis thaliana* (NCBI Identifier No. gi 3608136) and *Pisum sativum* (NCBI Identifier No. gi 3869255). In the process of comparing the soybean clones it was found that they had overlapping regions of homology. Using this homology it was possible to align all the soybean clones, including sel.pk0035.e10, in order to assemble a contig encoding a unique soybean DAD1 protein. The assembled contig revealed similarity of the encoded protein to

35 DAD1 from *Pisum sativum* (NCBI Identifier No. gi 3869255). The BLASTX search using EST sequence from a corn clone, cbn10.pk0058.g12, revealed similarity of the protein encoded by the cDNA to DAD1 from *Oryza sativa* (NCBI Identifier No. gi 2723473). The BLAST results for the corn and *Phaseolus* ESTs and the soybean contig are shown in Table 3:

TABLE 3
BLAST Results for Clones Encoding Polypeptides Homologous
to *Arabidopsis thaliana* DAD1 Proteins

Clone	BLAST pLog Score
plht.pk0012.c8	64.00
cbn10.pk0058.g12	67.30
Contig composed of:	62.00
se1.pk0035.e10	
ses2w.pk0011.h3	
sdp4c.pk007.e17	
sl1.pk0106.d1	
sls1c.pk010.h9	

5

The sequence of the cDNA insert in clone plht.pk0012.c8 encodes 100% of a *Phaseolus* DAD1 protein and is shown in SEQ ID NO:1; the deduced amino acid sequence of this cDNA is shown in SEQ ID NO:2. The amino acid sequence set forth in SEQ ID NO:2 was evaluated by BLASTP, yielding a pLog value of 53.00 versus an *Arabidopsis* *thaliana* (NCBI Identifier No. gi 2623638) DAD1 sequence. A calculation of the percent similarity of the amino acid sequence set forth in SEQ ID NO:2 and the *Arabidopsis thaliana* sequence can be seen in Table 4.

10

The sequence of the entire cDNA insert in clone cbn10.pk0058.g12 was determined and is shown in SEQ ID NO:3; the deduced amino acid sequence of this cDNA is shown in SEQ ID NO:4. The amino acid sequence set forth in SEQ ID NO:4 was evaluated by BLASTP, yielding a pLog value of 57.00 versus the *Oryza sativa* (NCBI Identifier No. gi 2723473) sequence. A calculation of the percent similarity of the amino acid sequence set forth in SEQ ID NO:4 and the *Oryza sativa* sequence can be seen in Table 4.

15

The sequence of the entire contig composed of clones se1.pk0035.e10, ses2w.pk0011.h3, sdp4c.pk007.e17, sl1.pk0106.d1 and sls1c.pk010.h9 is shown in SEQ ID NO:5; the deduced amino acid sequence of this cDNA is shown in SEQ ID NO:6. The amino acid sequence set forth in SEQ ID NO:6 was evaluated by BLASTP, yielding a pLog value of 52.52 versus the *Pisum sativum* (NCBI Identifier No. gi 3869255) sequence. A calculation of the percent similarity of the amino acid sequence set forth in SEQ ID NO:6 and the *Pisum sativum* sequence can be seen in Table 4. Figure 1 presents an alignment of the amino acid sequences set forth in SEQ ID NOs:2, 4 and 6 and the *Arabidopsis thaliana*, *Oryza sativa* and *Pisum sativum* sequences.

20

25

TABLE 4

Percent Identity of Amino Acid Sequences Deduced From the Nucleotide Sequences
of cDNA Clones Encoding Polypeptides Homologous
to *Arabidopsis thaliana*, *Oryza sativa* and *Pisum sativum* DAD1 Proteins

Clone	SEQ ID NO.	Percent Similarity to (Species)
plht.pk0012.c8	2	90% (<i>Arabidopsis thaliana</i> gi 2623638)
cbn10.pk0058.g12	4	97% (<i>Oryza sativa</i> gi 2723473)
A Contig composed of: sel.pk0035.e10 ses2w.pk0011.h3 sdp4c.pk007.e17 sl1.pk0106.d1 sls1c.pk010.h9	6	85% (<i>Pisum sativum</i> , gi 3869255)

- 5 Sequence alignments and percent similarity calculations were performed by the Clustal Algorithm (Higgins, D. G. et al., (1989) *CABIOS* 5(2):151-153), using the Megalign program of the LASARGENE bioinformatics computing suite (DNASTAR Inc., Madison, WI). Default parameters for the Clustal method for protein multiple alignments are: GAP PENALTY=10, GAP LENGTH PENALTY=10; for pairwise alignments KTUPLE 1, GAP
- 10 PENALTY=3, WINDOW=5 and DIAGONALS SAVED=5. Sequence alignments and BLAST scores and probabilities indicate that the instant nucleic acid fragments encode entire DAD1 proteins. These sequences represent the first corn, *Phaseolus* and soybean sequences encoding DAD1 proteins.

EXAMPLE 4

15 Characterization of cDNA Clones Encoding AAC-11 Homologs

- The BLASTX search using the EST sequences from clones p0031.ccmba65r, p0037.crwaj44r, p0049.curar45r, p0004.cb1ha38r, p0011.cdapf47r, cbn10.pk0041.h3, cen1.pk0032.d4, p0116.cesag38r, rls24.pk0025.b4, sdp2c.pk027.j3 and src3c.pk001.j10 revealed similarity of the proteins encoded by the cDNAs to AAC-11 from *Homo sapiens*
- 20 (NCBI Identifier No. gi 2623761). In the process of comparing the ESTs it was found that corn clones p0031.ccmba65r, p0037.crwaj44r, p0049.curar45r, p0004.cb1ha38r, p0011.cdapf47r, cbn10.pk0041.h3, cen1.pk0032.d4 and p0116.cesag38r had overlapping regions of homology. Soybean clones sdp2c.pk027.j3 and src3c.pk001.j10 were also found to have overlapping regions of homology. Using this homology it was possible to align the
- 25 ESTs and assemble two contigs encoding unique corn and soybean AAC-11 proteins. The BLAST results for each of the contigs and the rice EST are shown in Table 5:

TABLE 5

BLAST Results for Clones Encoding Polypeptides Homologous
to *Homo sapiens* AAC-11 Protein

Clone	BLAST pLog Score
Contig composed of: p0031.ccmba65r p0037.crwaj44r p0049.curar45r p0004.cb1ha38r p0011.cdapf47r cbn10.pk0041.h3 cen1.pk0032.d4 p0116.cesag38r	37.70
rls24.pk0025.b4	5.22
Contig composed of: sdp2c.pk027.j3 src3c.pk001.j10	16.70

5 The sequence of the corn contig composed of clones p0031.ccmba65r,
p0037.crwaj44r, p0049.curar45r, p0004.cb1ha38r, p0011.cdapf47r, cbn10.pk0041.h3,
cen1.pk0032.d4 and p0116.cesag38r encodes an entire AAC-11 protein and is shown in SEQ
ID NO:7; the deduced amino acid sequence of this contig is shown in SEQ ID NO:8. The
amino acid sequence set forth in SEQ ID NO:8 was evaluated by BLASTP, yielding a pLog
10 value of 49.70 versus the human sequence. A calculation of the percent similarity of the
amino acid sequence set forth in SEQ ID NO:8 and the *Homo sapiens* AAC-11 sequence
reveals that the corn AAC-11 amino acid sequence is 21% similar to the human AAC-11
protein. Figure 2 presents an alignment of the amino acid sequences set forth in SEQ ID
NO:8 and the human AAC-11 sequence.

15 Sequence alignments and percent similarity calculations were performed by the Clustal
Algorithm (Higgins, D. G. et al., (1989) *CABIOS* 5(2):151-153), using the Megalign
program of the LASARGENE bioinformatics computing suite (DNASTAR Inc., Madison,
WI). Default parameters for the Clustal method for protein multiple alignments are: GAP
PENALTY=10, GAP LENGTH PENALTY=10; for pairwise alignments KTUPLE 1, GAP
20 PENALTY=3, WINDOW=5 and DIAGONALS SAVED=5. Sequence alignments and
BLAST scores and probabilities indicate that the instant nucleic acid fragments encode
entire DAD1 proteins.

The sequence of a portion of the cDNA insert from clone rls24.pk0025.b4 is shown in
SEQ ID NO:9; the deduced amino acid sequence of this cDNA is shown in SEQ ID NO:10.
25 The sequence of the soybean contig composed of clones sdp2c.pk027.j3 and src3c.pk001.j10
is shown in SEQ ID NO:11; the deduced amino acid sequence of this contig is shown in
SEQ ID NO:12.

BLAST scores and probabilities indicate that the instant nucleic acid fragments encode entire or portions of AAC-11 proteins. These sequences represent the first plant sequences encoding AAC-11 proteins.

EXAMPLE 5

Expression of Chimeric Genes in Monocot Cells

A chimeric gene comprising a cDNA encoding an apoptosis inhibitory protein in sense orientation with respect to the maize 27 kD zein promoter that is located 5' to the cDNA fragment, and the 10 kD zein 3' end that is located 3' to the cDNA fragment, can be constructed. The cDNA fragment of this gene may be generated by polymerase chain reaction (PCR) of the cDNA clone using appropriate oligonucleotide primers. Cloning sites (NcoI or SmaI) can be incorporated into the oligonucleotides to provide proper orientation of the DNA fragment when inserted into the digested vector pML103 as described below. Amplification is then performed in a standard PCR. The amplified DNA is then digested with restriction enzymes NcoI and SmaI and fractionated on an agarose gel. The appropriate band can be isolated from the gel and combined with a 4.9 kb NcoI-SmaI fragment of the plasmid pML103. Plasmid pML103 has been deposited under the terms of the Budapest Treaty at ATCC (American Type Culture Collection, 10801 University Blvd., Manassas, VA 20110-2209), and bears accession number ATCC 97366. The DNA segment from pML103 contains a 1.05 kb Sall-NcoI promoter fragment of the maize 27 kD zein gene and a 0.96 kb SmaI-Sall fragment from the 3' end of the maize 10 kD zein gene in the vector pGem9Zf(+) (Promega). Vector and insert DNA can be ligated at 15°C overnight, essentially as described (Maniatis). The ligated DNA may then be used to transform *E. coli* XL1-Blue (Epicurian Coli XL-1 Blue™; Stratagene). Bacterial transformants can be screened by restriction enzyme digestion of plasmid DNA and limited nucleotide sequence analysis using the dideoxy chain termination method (Sequenase™ DNA Sequencing Kit; U. S. Biochemical). The resulting plasmid construct would comprise a chimeric gene encoding, in the 5' to 3' direction, the maize 27 kD zein promoter, a cDNA fragment encoding an apoptosis inhibitory protein, and the 10 kD zein 3' region.

The chimeric gene described above can then be introduced into corn cells by the following procedure. Immature corn embryos can be dissected from developing caryopses derived from crosses of the inbred corn lines H99 and LH132. The embryos are isolated 10 to 11 days after pollination when they are 1.0 to 1.5 mm long. The embryos are then placed with the axis-side facing down and in contact with agarose-solidified N6 medium (Chu et al., (1975) *Sci. Sin. Peking* 18:659-668). The embryos are kept in the dark at 27°C. Friable embryogenic callus consisting of undifferentiated masses of cells with somatic proembryoids and embryoids borne on suspensor structures proliferates from the scutellum of these immature embryos. The embryogenic callus isolated from the primary explant can be cultured on N6 medium and sub-cultured on this medium every 2 to 3 weeks.

The plasmid, p35S/Ac (obtained from Dr. Peter Eckes, Hoechst Ag, Frankfurt, Germany) may be used in transformation experiments in order to provide for a selectable marker. This plasmid contains the *Pat* gene (see European Patent Publication 0 242 236) which encodes phosphinothricin acetyl transferase (PAT). The enzyme PAT confers resistance to herbicidal glutamine synthetase inhibitors such as phosphinothricin. The *pat* gene in p35S/Ac is under the control of the 35S promoter from Cauliflower Mosaic Virus (Odell et al. (1985) *Nature* 313:810-812) and the 3' region of the nopaline synthase gene from the T-DNA of the Ti plasmid of *Agrobacterium tumefaciens*.

The particle bombardment method (Klein et al., (1987) *Nature* 327:70-73) may be used to transfer genes to the callus culture cells. According to this method, gold particles (1 μ m in diameter) are coated with DNA using the following technique. Ten μ g of plasmid DNAs are added to 50 μ L of a suspension of gold particles (60 mg per mL). Calcium chloride (50 μ L of a 2.5 M solution) and spermidine free base (20 μ L of a 1.0 M solution) are added to the particles. The suspension is vortexed during the addition of these solutions. After 10 minutes, the tubes are briefly centrifuged (5 sec at 15,000 rpm) and the supernatant removed. The particles are resuspended in 200 μ L of absolute ethanol, centrifuged again and the supernatant removed. The ethanol rinse is performed again and the particles resuspended in a final volume of 30 μ L of ethanol. An aliquot (5 μ L) of the DNA-coated gold particles can be placed in the center of a Kapton™ flying disc (Bio-Rad Labs). The particles are then accelerated into the corn tissue with a Biolistic™ PDS-1000/He (Bio-Rad Instruments, Hercules CA), using a helium pressure of 1000 psi, a gap distance of 0.5 cm and a flying distance of 1.0 cm.

For bombardment, the embryogenic tissue is placed on filter paper over agarose-solidified N6 medium. The tissue is arranged as a thin lawn and covered a circular area of about 5 cm in diameter. The petri dish containing the tissue can be placed in the chamber of the PDS-1000/He approximately 8 cm from the stopping screen. The air in the chamber is then evacuated to a vacuum of 28 inches of Hg. The macrocarrier is accelerated with a helium shock wave using a rupture membrane that bursts when the He pressure in the shock tube reaches 1000 psi.

Seven days after bombardment the tissue can be transferred to N6 medium that contains glufosinate (2 mg per liter) and lacks casein or proline. The tissue continues to grow slowly on this medium. After an additional 2 weeks the tissue can be transferred to fresh N6 medium containing glufosinate. After 6 weeks, areas of about 1 cm in diameter of actively growing callus can be identified on some of the plates containing the glufosinate-supplemented medium. These calli may continue to grow when sub-cultured on the selective medium.

Plants can be regenerated from the transgenic callus by first transferring clusters of tissue to N6 medium supplemented with 0.2 mg per liter of 2,4-D. After two weeks the

tissue can be transferred to regeneration medium (Fromm et al., (1990) *Bio/Technology* 8:833-839).

EXAMPLE 6

Expression of Chimeric Genes in Dicot Cells

5 A seed-specific expression cassette composed of the promoter and transcription terminator from the gene encoding the β subunit of the seed storage protein phaseolin from the bean *Phaseolus vulgaris* (Doyle et al. (1986) *J. Biol. Chem.* 261:9228-9238) can be used for expression of the instant apoptosis inhibitory proteins in transformed soybean. The phaseolin cassette includes about 500 nucleotides upstream (5') from the translation initiation
10 codon and about 1650 nucleotides downstream (3') from the translation stop codon of phaseolin. Between the 5' and 3' regions are the unique restriction endonuclease sites Nco I (which includes the ATG translation initiation codon), Sma I, Kpn I and Xba I. The entire cassette is flanked by Hind III sites.

The cDNA fragment of this gene may be generated by polymerase chain reaction
15 (PCR) of the cDNA clone using appropriate oligonucleotide primers. Cloning sites can be incorporated into the oligonucleotides to provide proper orientation of the DNA fragment when inserted into the expression vector. Amplification is then performed as described above, and the isolated fragment is inserted into a pUC18 vector carrying the seed expression cassette.

20 Soybean embryos may then be transformed with the expression vector comprising sequences encoding apoptosis inhibitory proteins. To induce somatic embryos, cotyledons, 3-5 mm in length dissected from surface sterilized, immature seeds of the soybean cultivar A2872, can be cultured in the light or dark at 26°C on an appropriate agar medium for 6-10 weeks. Somatic embryos which produce secondary embryos are then excised and
25 placed into a suitable liquid medium. After repeated selection for clusters of somatic embryos which multiplied as early, globular staged embryos, the suspensions are maintained as described below.

Soybean embryogenic suspension cultures can maintained in 35 mL liquid media on a rotary shaker, 150 rpm, at 26°C with florescent lights on a 16:8 hour day/night schedule.

30 Cultures are subcultured every two weeks by inoculating approximately 35 mg of tissue into 35 mL of liquid medium.

Soybean embryogenic suspension cultures may then be transformed by the method of particle gun bombardment (Kline et al. (1987) *Nature* (London) 327:70, U.S. Patent No. 4,945,050). A DuPont Biolistic™ PDS1000/HE instrument (helium retrofit) can be used
35 for these transformations.

A selectable marker gene which can be used to facilitate soybean transformation is a chimeric gene composed of the 35S promoter from Cauliflower Mosaic Virus (Odell et al.(1985) *Nature* 313:810-812), the hygromycin phosphotransferase gene from plasmid pJR225 (from *E. coli*; Gritz et al.(1983) *Gene* 25:179-188) and the 3' region of the nopaline

synthase gene from the T-DNA of the Ti plasmid of *Agrobacterium tumefaciens*. The seed expression cassette comprising the phaseolin 5' region, the fragment encoding the apoptosis inhibitory protein and the phaseolin 3' region can be isolated as a restriction fragment. This fragment can then be inserted into a unique restriction site of the vector carrying the marker gene.

To 50 μ L of a 60 mg/mL 1 μ m gold particle suspension is added (in order): 5 μ L DNA (1 μ g/ μ L), 20 μ L spermidine (0.1 M), and 50 μ L CaCl_2 (2.5 M). The particle preparation is then agitated for three minutes, spun in a microfuge for 10 seconds and the supernatant removed. The DNA-coated particles are then washed once in 400 μ L 70% ethanol and resuspended in 40 μ L of anhydrous ethanol. The DNA/particle suspension can be sonicated three times for one second each. Five μ L of the DNA-coated gold particles are then loaded on each macro carrier disk.

Approximately 300-400 mg of a two-week-old suspension culture is placed in an empty 60x15 mm petri dish and the residual liquid removed from the tissue with a pipette. For each transformation experiment, approximately 5-10 plates of tissue are normally bombarded. Membrane rupture pressure is set at 1100 psi and the chamber is evacuated to a vacuum of 28 inches mercury. The tissue is placed approximately 3.5 inches away from the retaining screen and bombarded three times. Following bombardment, the tissue can be divided in half and placed back into liquid and cultured as described above.

Five to seven days post bombardment, the liquid media may be exchanged with fresh media, and eleven to twelve days post bombardment with fresh media containing 50 mg/mL hygromycin. This selective media can be refreshed weekly. Seven to eight weeks post bombardment, green, transformed tissue may be observed growing from untransformed, necrotic embryogenic clusters. Isolated green tissue is removed and inoculated into individual flasks to generate new, clonally propagated, transformed embryogenic suspension cultures. Each new line may be treated as an independent transformation event. These suspensions can then be subcultured and maintained as clusters of immature embryos or regenerated into whole plants by maturation and germination of individual somatic embryos.

EXAMPLE 7

Expression of Chimeric Genes in Microbial Cells

The cDNAs encoding the instant apoptosis inhibitory proteins can be inserted into the T7 *E. coli* expression vector pBT430. This vector is a derivative of pET-3a (Rosenberg et al. (1987) *Gene* 56:125-135) which employs the bacteriophage T7 RNA polymerase/T7 promoter system. Plasmid pBT430 was constructed by first destroying the EcoR I and Hind III sites in pET-3a at their original positions. An oligonucleotide adaptor containing EcoR I and Hind III sites was inserted at the BamH I site of pET-3a. This created pET-3aM with additional unique cloning sites for insertion of genes into the expression vector. Then, the Nde I site at the position of translation initiation was converted to an Nco I site using

oligonucleotide-directed mutagenesis. The DNA sequence of pET-3aM in this region, 5'-CATATGG, was converted to 5'-CCCATGG in pBT430.

5 Plasmid DNA containing a cDNA may be appropriately digested to release a nucleic acid fragment encoding the protein. This fragment may then be purified on a 1% NuSieve GTG™ low melting agarose gel (FMC). Buffer and agarose contain 10 µg/ml ethidium bromide for visualization of the DNA fragment. The fragment can then be purified from the agarose gel by digestion with GELase™ (Epicentre Technologies) according to the manufacturer's instructions, ethanol precipitated, dried and resuspended in 20 µL of water. Appropriate oligonucleotide adapters may be ligated to the fragment using T4 DNA ligase 10 (New England Biolabs, Beverly, MA). The fragment containing the ligated adapters can be purified from the excess adapters using low melting agarose as described above. The vector pBT430 is digested, dephosphorylated with alkaline phosphatase (NEB) and deproteinized with phenol/chloroform as described above. The prepared vector pBT430 and fragment can then be ligated at 16°C for 15 hours followed by transformation into DH5 electrocompetent 15 cells (GIBCO BRL). Transformants can be selected on agar plates containing LB media and 100 µg/mL ampicillin. Transformants containing the gene encoding the apoptosis inhibitory protein are then screened for the correct orientation with respect to the T7 promoter by restriction enzyme analysis.

For high level expression, a plasmid clone with the cDNA insert in the correct 20 orientation relative to the T7 promoter can be transformed into *E. coli* strain BL21(DE3) (Studier et al. (1986) *J. Mol. Biol.* 189:113-130). Cultures are grown in LB medium containing ampicillin (100 mg/L) at 25°C. At an optical density at 600 nm of approximately 1, IPTG (isopropylthio-β-galactoside, the inducer) can be added to a final concentration of 0.4 mM and incubation can be continued for 3 h at 25°. Cells are then harvested by 25 centrifugation and re-suspended in 50 µL of 50 mM Tris-HCl at pH 8.0 containing 0.1 mM DTT and 0.2 mM phenyl methylsulfonyl fluoride. A small amount of 1 mm glass beads can be added and the mixture sonicated 3 times for about 5 seconds each time with a microprobe sonicator. The mixture is centrifuged and the protein concentration of the supernatant determined. One µg of protein from the soluble fraction of the culture can be separated by 30 SDS-polyacrylamide gel electrophoresis. Gels can be observed for protein bands migrating at the expected molecular weight.

CLAIMS

What is claimed is:

1. An isolated nucleic acid fragment encoding all or a substantial portion of a DAD1 protein comprising a member selected from the group consisting of:
 - 5 (a) an isolated nucleic acid fragment encoding all or a substantial portion of the amino acid sequence set forth in a member selected from the group consisting of SEQ ID NO:2, 4 and 6;
 - (b) an isolated nucleic acid fragment that is substantially similar to an isolated nucleic acid fragment encoding all or a substantial portion of
10 the amino acid sequence set forth in a member selected from the group consisting of SEQ ID NO:2, 4 and 6; and
 - (c) an isolated nucleic acid fragment that is complementary to (a) or (b).
2. The isolated nucleic acid fragment of Claim 1 wherein the nucleotide sequence of the fragment comprises all or a portion of the sequence set forth in a member selected
15 from the group consisting of SEQ ID NO:1, 3 and 5.
3. The isolated nucleic acid fragment of Claim 1 wherein the isolated nucleic acid fragment hybridizes under stringent conditions to an isolated nucleic acid fragment encoding all or a substantial portion of the amino acid sequence set forth in SEQ ID NO:2.
4. The isolated nucleic acid fragment of Claim 1 wherein the isolated nucleic acid
20 fragment hybridizes under stringent conditions to an isolated nucleic acid fragment encoding all or a substantial portion of the amino acid sequence set forth in SEQ ID NO:4.
5. The isolated nucleic acid fragment of Claim 1 wherein the isolated nucleic acid fragment hybridizes under stringent conditions to an isolated nucleic acid fragment encoding all or a substantial portion of the amino acid sequence set forth in SEQ ID NO:5.
- 25 6. The isolated nucleic acid fragment of Claim 1 wherein the isolated nucleic acid fragment is at least 85% similar to an isolated nucleic acid fragment encoding all or a substantial portion of the amino acid sequence set forth in SEQ ID NO:2.
7. The isolated nucleic acid fragment of Claim 1 wherein the isolated nucleic acid fragment is at least 85% similar to an isolated nucleic acid fragment encoding all or a
30 substantial portion of the amino acid sequence set forth in SEQ ID NO:4.
8. The isolated nucleic acid fragment of Claim 1 wherein the isolated nucleic acid fragment is at least 85% similar to an isolated nucleic acid fragment encoding all or a substantial portion of the amino acid sequence set forth in SEQ ID NO:6.
9. A chimeric gene comprising the nucleic acid fragment of Claim 1 operably
35 linked to suitable regulatory sequences.
10. A transformed host cell comprising the chimeric gene of Claim 9.
11. A DAD1 polypeptide comprising all or a substantial portion of the amino acid sequence set forth in a member selected from the group consisting of SEQ ID NO:2, 4 and 6.

12. An isolated nucleic acid fragment encoding all or a substantial portion of an AAC-11 protein comprising a member selected from the group consisting of:

(a) an isolated nucleic acid fragment encoding all or a substantial portion of the amino acid sequence set forth in a member selected from the group consisting of SEQ ID NO:8, 10 and 12;

(b) an isolated nucleic acid fragment that is substantially similar to an isolated nucleic acid fragment encoding all or a substantial portion of the amino acid sequence set forth in a member selected from the group consisting of SEQ ID NO:8, 10 and 12; and

(c) an isolated nucleic acid fragment that is complementary to (a) or (b).

13. The isolated nucleic acid fragment of Claim 12 wherein the nucleotide sequence of the fragment comprises all or a portion of the sequence set forth in a member selected from the group consisting of SEQ ID NO:7, 9 and 11.

14. The isolated nucleic acid fragment of Claim 12 wherein the isolated nucleic acid fragment hybridizes under stringent conditions to an isolated nucleic acid fragment encoding all or a substantial portion of the amino acid sequence set forth in SEQ ID NO:8.

15. The isolated nucleic acid fragment of Claim 12 wherein the isolated nucleic acid fragment hybridizes under stringent conditions to an isolated nucleic acid fragment encoding all or a substantial portion of the amino acid sequence set forth in SEQ ID NO:10.

16. The isolated nucleic acid fragment of Claim 12 wherein the isolated nucleic acid fragment hybridizes under stringent conditions to an isolated nucleic acid fragment encoding all or a substantial portion of the amino acid sequence set forth in SEQ ID NO:12.

17. The isolated nucleic acid fragment of Claim 12 wherein the isolated nucleic acid fragment is at least 85% similar to an isolated nucleic acid fragment encoding all or a substantial portion of the amino acid sequence set forth in SEQ ID NO:8.

18. The isolated nucleic acid fragment of Claim 12 wherein the isolated nucleic acid fragment is at least 85% similar to an isolated nucleic acid fragment encoding all or a substantial portion of the amino acid sequence set forth in SEQ ID NO:10.

19. The isolated nucleic acid fragment of Claim 12 wherein the isolated nucleic acid fragment is at least 85% similar to an isolated nucleic acid fragment encoding all or a substantial portion of the amino acid sequence set forth in SEQ ID NO:12.

20. A chimeric gene comprising the nucleic acid fragment of Claim 12 operably linked to suitable regulatory sequences.

21. A transformed host cell comprising the chimeric gene of Claim 20.

22. An AAC-11 polypeptide comprising all or a substantial portion of the amino acid sequence set forth in a member selected from the group consisting of SEQ ID NO:8, 10 and 12.

23. A method of altering the level of expression of an apoptosis inhibitory protein in a host cell comprising:

- (a) transforming a host cell with the chimeric gene of any of Claims 9 and 20; and
- (b) growing the transformed host cell produced in step (a) under conditions that are suitable for expression of the chimeric gene

5 wherein expression of the chimeric gene results in production of altered levels of an apoptosis inhibitory protein in the transformed host cell.

24. A method of obtaining a nucleic acid fragment encoding all or a substantial portion of the amino acid sequence encoding an apoptosis inhibitory protein comprising:

- 10 (a) probing a cDNA or genomic library with the nucleic acid fragment of any of Claims 1 and 12 ;
- (b) identifying a DNA clone that hybridizes with the nucleic acid fragment of any of Claims 1 and 12;
- (c) isolating the DNA clone identified in step (b); and
- 15 (d) sequencing the cDNA or genomic fragment that comprises the clone isolated in step (c)

wherein the sequenced nucleic acid fragment encodes all or a substantial portion of the amino acid sequence encoding an apoptosis inhibitory protein.

25. A method of obtaining a nucleic acid fragment encoding a substantial portion of an amino acid sequence encoding an apoptosis inhibitory protein comprising:

- 20 (a) synthesizing an oligonucleotide primer corresponding to a portion of the sequence set forth in any of SEQ ID NOs:1, 3, 5, 7, 9 and 11 ; and
- (b) amplifying a cDNA insert present in a cloning vector using the oligonucleotide primer of step (a) and a primer representing sequences of the cloning vector

25 wherein the amplified nucleic acid fragment encodes a substantial portion of an amino acid sequence encoding an apoptosis inhibitory protein.

26. The product of the method of Claim 24.

27. The product of the method of Claim 25.

1	60		
		MV-----KSTSKDAQDLFHSLSAY-TATPTNLKIIDLYVCFVAVFTALIQVAYMALVGS	
		MP-----RATSDAKLLIQSLGKAY-AATPTNLKIIDLYVVFVAVATALIQQVVMGIVGS	
		MA-----KTSSTTKDAQDLFHAWSAY-SATPTNLKIIDLYVVFVAVTALLQDVYMALVGP	
		MA-----RSSSKDAQDLFRALWSAY-AATPTNLKIIDLYVMFAVFTALIQVVMALVGS	
		MP-----RATSDAKLLIQSLGKAY-AATPTNLKIIDLYVGFVAVATALIQQVAYMGLVGS	
		MA-----PRSSSKDAQDLFRALWSAY-AATPTNLKIIDLYVIYAVFTAFIQVVMALVGS	
SEQ ID NO: 13			
SEQ ID NO: 14			
SEQ ID NO: 15			
SEQ ID NO: 2			
SEQ ID NO: 4			
SEQ ID NO: 6			
	120		
		FPFNSFLSGVLSGIGTAVLAVCLRIQVNKENK-EFKDLAPERAFADFVLCNLVHLVIMN	
		FPFNSFLSGVLSGIGTAVLAVCLRIQVNKDNK-EFKDLPPERAFADFVLCNLVHLVIMN	
		FPFNSFLSGVLSGIGTAVLAVCLRIQVNKENK-EFKDLGPERAFADFVLCNLVHLVIMN	
		FPFNSFLSGVLSGIGTAVLAVCLRIQVNKENK-EFKDLAPERAFADFVLCNLVHLVIMN	
		FPFNSFLSGVLSGIGTAVLAVCLRIQVNKDNK-EFKDLPPERAFADFVLCNLVHLVIMN	
		FPFNSFLSGVLSGIGTAVLAVCLRIQVNKENK-EFKDLAPERAFADFVLCNLVHLVIMN	
SEQ ID NO: 13			
SEQ ID NO: 14			
SEQ ID NO: 15			
SEQ ID NO: 2			
SEQ ID NO: 4			
SEQ ID NO: 6			
	121		
		FL-G	
		FL-G	
		FLG.	
		FLG.	
		FLG.	
		FLG.	
SEQ ID NO: 13			
SEQ ID NO: 14			
SEQ ID NO: 15			
SEQ ID NO: 2			
SEQ ID NO: 4			
SEQ ID NO: 6			

FIG. 1

2/3

1	60	
		MGLSLT--MPTVEELYRNYGILADATEQVGQHKDAYQVILDGVKG-GTKEKRLLAAQFIPK
SEQ ID NO:16		
SEQ ID NO:8		MAAAAADDAEAEVERLY-ELGERLSSANDKSEHAADYEAI IAAVKGSAAKAKQLAAQLIPR
	120	
61		FFKHFPPELADSAINAQDLCEDEDVSIRROAIKELPOF--ATGENLPRVADILTQLLQTD
SEQ ID NO:16		
SEQ ID NO:8		FFRSFPALGTRAMSAMFDLVDMEELAIRIQAIRGFPLLGKDT-EFVSKIADVLGQLLTSE
	180	
121		DSAEFNLVNNALLSIFKMDAKGTLGGGLFSQIL---QGEDIVRERAIFLSTKLKTLDPDEV
SEQ ID NO:16		
SEQ ID NO:8		ENVERDAVHKALMSLIRQDVKNLSQPLFKHVE----QGSEIREKIIICFLRDKVFLKAEAL
	240	
181		LT--KEVEELILTESKKVLEDVTGEEFVLFMKILSGLSLQTVSGR---QQLVELVAEQA
SEQ ID NO:16		
SEQ ID NO:8		LKPQAEEMERFITDLIKKSVQDVTGSEFEFELFMGFLRXWSIFGDSAPRESFQELIEIIQAQA
	300	
241		DLEQTFNPSPDCVDRLLQCTRQAVPLFSKNVHSTRFVTFCEQVLPNLGTLTTPVEGLD
SEQ ID NO:16		
SEQ ID NO:8		DLNSQFNVSDDIDHIERWISCMYMALPIFMRGASASKFLNYFVKQIVPAFEKIPE-----E
	360	
301		IQLEVLKLLAEMSSFCGDMEKLETNLRKLFDKLLEYMPLPPEEAENGENAGNEEPKLQFS
SEQ ID NO:16		
SEQ ID NO:8		KKLDLLKTIASSSPYATAQDSRQL-LPSVVQLLNKYMP-----GKKV-DD---INHN

FIG. 2

3/3

SEQ ID NO:16	361	420
SEQ ID NO:8	YVECLLYSEHQGRKLPDELTA-----KL-----NAEKLHES-----KIRLOQFARGLO	
	YVECLLYTYHHLAHKTPNTNTNSLCGYKIVTGQPSDRLGEDFTEHYKDFTERLTGTTEETVR	
SEQ ID NO:16	421	480
SEQ ID NO:8	VYIRQLRLALQG--KTGEALKTEENKIKV-----VALKITNNINVLIKDLFHIPPSY	
	AASKRLTQGMADFSKAISSAKTEEEKTKIKGDQQTSTRTRMSYNNILAMTQSLHKSPLF	
SEQ ID NO:16	481	540
SEQ ID NO:8	--KSTVTLSW-----KPVQKVEIGQKRASEDTTSGSPPKSSAGPKRMPGRFITLPVGNIA	
	IGDKKITLSWMEQPNKAAATKAGGKRSQPSPTNGNDPANKKGRGG--MQNQLVNRAFEGLS	
SEQ ID NO:16	541	588
SEQ ID NO:8	AIWATLIMRGAFRGSK-----WPRLGH--TKEIVVGERLY	
	HVGRGSGRXRGXGGPRKRKRMEWGTTEMTWKLCNLTRDELSASTRE.	

FIG. 2 (Continued)

SEQUENCE LISTING

<110> E. I. DU PONT DE NEMOURS AND COMPANY
 <120> INHIBITORS OF APOPTOSIS PROTEINS IN PLANTS
 <130> BB-1148
 <150> 60/078,144
 <151> MARCH 16, 1998
 <160> 16
 <170> MICROSOFT WORD VERSION 7.0A
 <210> 1
 <211> 499
 <212> DNA
 <213> Phaseolus lunatus

<400> 1
 gcgaagatct gagagaatgg cgaggtctag cagcaaggat gcacaagacc ttttccgagc 60
 tcttttggct gcttatgccc caacccccac aaatctcaag atcattgacc tctacgtcat 120
 gttcgcgtgt ttcaccgctc tcatccaggt agtttacctg gctttgggtgg gatcatttcc 180
 ttttaactcc ttcctatcag gagtactttc ttgtgtcggg actgctgttc ttgctgtttg 240
 tctcaggatc caagtgaata aagagaataa ggaattcaag gatcttgac ctgagcgagc 300
 ttttgcagat tttgttctct gtaatctggt gcttcatttg gtgatcatga acttccttgg 360
 ttaatttgag ttcattgtggc tggtgttggt tttgatcaaa ccttggataa taaaaagtaa 420
 tagtagtata cctagacttt tgtaatagta tttatagaca gtaacttcca actaactgct 480
 ttagtatttt gttgattcc 499

<210> 2
 <211> 115
 <212> PRT
 <213> Phaseolus lunatus

<400> 2
 Met Ala Arg Ser Ser Ser Lys Asp Ala Gln Asp Leu Phe Arg Ala Leu
 1 5 10 15
 Trp Ser Ala Tyr Ala Ala Thr Pro Thr Asn Leu Lys Ile Ile Asp Leu
 20 25 30
 Tyr Val Met Phe Ala Val Phe Thr Ala Leu Ile Gln Val Val Tyr Met
 35 40 45
 Ala Leu Val Gly Ser Phe Pro Phe Asn Ser Phe Leu Ser Gly Val Leu
 50 55 60
 Ser Cys Val Gly Thr Ala Val Leu Ala Val Cys Leu Arg Ile Gln Val
 65 70 75 80
 Asn Lys Glu Asn Lys Glu Phe Lys Asp Leu Ala Pro Glu Arg Ala Phe
 85 90 95
 Ala Asp Phe Val Leu Cys Asn Leu Val Leu His Leu Val Ile Met Asn
 100 105 110
 Phe Leu Gly
 115

<210> 3
 <211> 712
 <212> DNA
 <213> Zea mays

```

<400> 3
gcacgagggc cgctgccga ccccgacgcc tgctgccag gtcttcgccg gcgacgagca 60
ctccaccaga cgagagggga ttccaagatg ccgagggcca ccagcgacgc gaagctcctg 120
atccagtccc tcggcaaggc gtacgtgccc acaccaacaa atctcaagat tattgacctc 180
tacgtgggtt ttgcggttgc cactgccctt attcaggttg cttacatggg attggttggg 240
tcgtttccct tcaactcctt cctctcagga gtcctttcat gcataggaac tgcagttctt 300
gctgtttgcc tccgcattca agtgaacaaa gacaacaaag aattcaagga ccttccccca 360
gaaagggcct ttgctgattt cgtcctatgc aatctggtgc tccacctggg gatcatgaat 420
ttcctcggat aagcaactgc tgcaccatgt tggttaaagg ttttgtagcc ccagggttgg 480
gtcgtcgatt gttgccttta aatgtttgga actgttgga tcgtgatgtc gaatatccat 540
atgatctgtt gaaggattac ttgtgtaagc tgagtattcc cggaggggaac tattagtcga 600
atggacagtt tgcacagcgc tgagaatgtg acctagcatg ttctttattt gaagaagata 660
taattcattt ttcaaaaaaa aaaaaaaaaa aactcgaggg gggcccgtac cc 712

```

```

<210> 4
<211> 114
<212> PRT
<213> Zea mays

```

```

<400> 4
Met Pro Arg Ala Thr Ser Asp Ala Lys Leu Leu Ile Gln Ser Leu Gly
1 5 10 15
Lys Ala Tyr Ala Ala Thr Pro Thr Asn Leu Lys Ile Ile Asp Leu Tyr
20 25 30
Val Gly Phe Ala Val Ala Thr Ala Leu Ile Gln Val Ala Tyr Met Gly
35 40 45
Leu Val Gly Ser Phe Pro Phe Asn Ser Phe Leu Ser Gly Val Leu Ser
50 55 60
Cys Ile Gly Thr Ala Val Leu Ala Val Cys Leu Arg Ile Gln Val Asn
65 70 75 80
Lys Asp Asn Lys Glu Phe Lys Asp Leu Pro Pro Glu Arg Ala Phe Ala
85 90 95
Asp Phe Val Leu Cys Asn Leu Val Leu His Leu Val Ile Met Asn Phe
100 105 110

```

Leu Gly

```

<210> 5
<211> 917
<212> DNA
<213> Glycine max

```

```

<220>
<221> unsure
<222> (631)

```

```

<220>
<221> unsure
<222> (636)

```

```

<220>
<221> unsure
<222> (652)

```

```

<220>
<221> unsure
<222> (657)

```

<220>
 <221> unsure
 <222> (719)

 <220>
 <221> unsure
 <222> (785)

 <220>
 <221> unsure
 <222> (843)

 <220>
 <221> unsure
 <222> (849)

 <220>
 <221> unsure
 <222> (859) .. (860)

 <220>
 <221> unsure
 <222> (886)

 <220>
 <221> unsure
 <222> (890)

 <220>
 <221> unsure
 <222> (897)

 <220>
 <221> unsure
 <222> (906)

<400> 5
 ggcaaatcgc gaagactaga tctgatctga gagaatggct cctcgggtcta gcagcaagga 60
 cgcccaagac cttttccgcg ctctttgggtc tgcttatgct gcaacccccca ctaatctcaa 120
 gatcattgat ctctatgtca tctatgccgt attcaccgct ttcattccagg ttgtttacat 180
 ggctttgggtt ggatcatttc catttaactc cttcctatca ggagtacttt cttgtgtagg 240
 aactgctgtt cttgctgttt gtctcaggat ccaagtgaat aaagagaata aggaattcaa 300
 ggatcttgca cctgagcgcg cttttgcgga ttttgttctc tgtaatttgg tgcttcattt 360
 ggtgatcatg aacttccttg gttaaattgg gtttgtgtgg cggttgttgt ttctgattga 420
 acccttcgat aataaaaatt aaatagtagt atacctagac ttttgtaata gtatttatag 480
 acagtagccg gcattctact aattgcttta gctttatgtt gattaccccc acctccatt 540
 tttgggttcc ctgttttgaa cgaagagatt ttgccatctt ttgaagttaa aagtactttt 600
 gaatggcgaa ataaagaagg attgttatta naaanaaaaa aaaaataaca anatatnaac 660
 gcttacattt aagtggcact ttcggggaaa tgtgcgcgga accctattgt taatttccna 720
 aatacatcaa atagtaccgc caagaacata accctgataa agctcataat atgaaaagga 780
 gatanatatc acattcgtgt cgcctaacct ttttgcggat ttgcctcccg gttttgccaa 840
 canaaccng gaaataaann gcgaaatact ggtgccaatg gtaacnatgn ttcaaangga 900
 aatctnaaat tcccaag 917

<210> 6
 <211> 116
 <212> PRT
 <213> Glycine max

<400> 6
 Met Ala Pro Arg Ser Ser Ser Lys Asp Ala Gln Asp Leu Phe Arg Ala
 1 5 10 15

Leu Trp Ser Ala Tyr Ala Ala Thr Pro Thr Asn Leu Lys Ile Ile Asp
 20 25 30
 Leu Tyr Val Ile Tyr Ala Val Phe Thr Ala Phe Ile Gln Val Val Tyr
 35 40 45
 Met Ala Leu Val Gly Ser Phe Pro Phe Asn Ser Phe Leu Ser Gly Val
 50 55 60
 Leu Ser Cys Val Gly Thr Ala Val Leu Ala Val Cys Leu Arg Ile Gln
 65 70 75 80
 Val Asn Lys Glu Asn Lys Glu Phe Lys Asp Leu Ala Pro Glu Arg Ala
 85 90 95
 Phe Ala Asp Phe Val Leu Cys Asn Leu Val Leu His Leu Val Ile Met
 100 105 110
 Asn Phe Leu Gly
 115

<210> 7
 <211> 2671
 <212> DNA
 <213> Zea mays

<220>
 <221> unsure
 <222> (690)

<220>
 <221> unsure
 <222> (2161)

<220>
 <221> unsure
 <222> (2180)

<400> 7
 gcatcccttc ccagttctgt cctctcgaac ccctaactcc aaaaaccctc gctctcctct 60
 catggccgcc gccgcagctg acgacgcagc cgaggtggag cggctttacg agctcggcga 120
 gcgcctctcc tccgccaatg acaagtccga gcatgcggcg gactacgagg cgattattgc 180
 agcgggtgaag ggacagagtg ccaaggcgaa gcagctcgca gcgcagctta tccccaggtt 240
 cttccggagc ttccctgcac tcggcacgcg cgccatgtca gccatgttcg atctcgtcga 300
 tatggaggag ctgcgatca gaatacaagc tttcgtggc tttccacttc ttggcaaaga 360
 tactgaattt gtgtcaaaaa ttgcagatgt tttgggtcag ctccctacaa gcgaggaaaa 420
 tgttgagcgt gatgctgttc ataaagcgct catgtccctt atacggcaag atgttaaaaa 480
 ttcattacaa cctttattta agcatgtgga gcaaggatca gagattcgtg agaagattat 540
 ttgttttctt cgagacaagg tctttcctct taaagcagag ctgctgaaac ctcaagcaga 600
 aatggagaga tttataacgg atttgataaa gaaaagtgtg caagatgtaa ctgggttcaga 660
 attcgaacta ttcattgggt tcttgcgaaan ttggagcata tttggggatt ctgctcctag 720
 agagtccttt caagaactaa ttgaaattat tcaagcacag gctgatctga attcacaatt 780
 caacgtttct gacattgacc acattgagag gtggatttca tgcattgata tggctcttcc 840
 gatcttcatt agaggagcat cagcaagcaa gttcctcaat tacttcgtta agcaaattgt 900
 tccagcattc gagaagattc ctgaagaaaa gaaactggat ttgctcaaga ctattgcttc 960
 aagttcaccg tacgcgacag ctcaagattc acgtcagctg cttccatctg ttgttcagtt 1020
 actcaacaaa tatatgcctg ggaagaaggt ggacgatatc aaccataatt atgttgaatg 1080
 cttgctgtac atttggtcga atttggtcca taagactcca aacacaacga acagtctatg 1140
 tggttacaag attgttactg ggcaaccatc ggatagactt ggagaggact tcacagagca 1200
 ttacaaagat ttacagaga ggtaaactgg aacagaagag acggtaagag cagcctcaaa 1260
 gcgactaact caggggaatg cagatttcag caaggcaata tcttcagcaa aaaccgaaga 1320
 ggaaaaaact aaaattaaag gcgatcaaca aacttcaaca aggacaatga ggtcatataa 1380
 caatatattg gcaatgacac agtcattgca ttcaaaatcc cttttattta tcggtgataa 1440
 gaaaatcact ctgtcatgga tggagcagcc caacaaagca gcagctacga aagcaggggg 1500
 gaagaggtca caacctagta caaatgggaa tgaccctgca aacaagaagg ggagaggagg 1560

```

aatgcaaaac cagctagtga acagagcttt tgaaggactg tctcatgttg gaagaggcag 1620
tggaagaggt cggggcaagg gtgggccgag gaagaggaag aggatggagt ggggtaccac 1680
tgagatgacc tggaaattat gttcaaacct gacaagggat gagctttctg cttccaccag 1740
agagtaaact ccaaggtcga tgttttattg gtgtgcatta ctgcaacgcg ttgaaatgga 1800
taaggcaca caacagcaga aatgcgtaca cagagaaagg atgctaagaa atatctgcaa 1860
gtttgtgcat ctttcttacc catttaccat ctcacgtgt tctttgccac cctaaccgtc 1920
gtgtcacctg cgttggctgg ctgtttgatg aactgggcag ttcgatatct ttgttcttta 1980
ttttatttta cagtgtttga agagacgacc aagcttgtgc tgactttgtt tgagttcggt 2040
ttatgtttcg tccttgact gacacagatg tattagtgt gtttaacttt tatgtaacga 2100
ttgattagct gtaataaag ttaagcattt aataagctat ttaaaaaaaa aaaaaaaacc 2160
nkrcgggatt ggagttgggn cccactgagg atgaccttgg aaatttattg ttcaaaacct 2220
tgaccaaggg gatgagcttt ctgcttccac cagagagtaa actccaaggt cgatgtttta 2280
tttgtgtgca ttactgcaac gcgttgaaat ggatcaaggc acacaacagc agaaatgcgt 2340
acacagagaa aggatgctaa gaaatatctg caagtttgtg catctttctt atccatttac 2400
catctcatcg tgttctttgc caccctaacc gtcgtgtcac ctgcttggc tggctgtttg 2460
atgaactggg cagttcgata tctttgttct ttattttatt ttacagtgtt tgaagagacg 2520
accaagcttg tgctgacttt gtttgagttc gttttatgtt tcgtccttgt actgacacag 2580
atgtattagt gatgtttaac ttttatgtaa cgattgatta gctgtaatat aagttaagca 2640
tttaataaagc tatttaaaaa aaaaaaaaaa a 2671

```

<210> 8
 <211> 561
 <212> PRT
 <213> Zea mays

<220>
 <221> UNSURE
 <222> (210)

<220>
 <221> UNSURE
 <222> (523)

<220>
 <221> UNSURE
 <222> (526)

<400> 8
 Met Ala Ala Ala Ala Ala Asp Asp Ala Ala Glu Val Glu Arg Leu Tyr
 1 5 10 15
 Glu Leu Gly Glu Arg Leu Ser Ser Ala Asn Asp Lys Ser Glu His Ala
 20 25 30
 Ala Asp Tyr Glu Ala Ile Ile Ala Ala Val Lys Gly Gln Ser Ala Lys
 35 40 45
 Ala Lys Gln Leu Ala Ala Gln Leu Ile Pro Arg Phe Phe Arg Ser Phe
 50 55 60
 Pro Ala Leu Gly Thr Arg Ala Met Ser Ala Met Phe Asp Leu Val Asp
 65 70 75 80
 Met Glu Glu Leu Ala Ile Arg Ile Gln Ala Ile Arg Gly Phe Pro Leu
 85 90 95
 Leu Gly Lys Asp Thr Glu Phe Val Ser Lys Ile Ala Asp Val Leu Gly
 100 105 110
 Gln Leu Leu Thr Ser Glu Glu Asn Val Glu Arg Asp Ala Val His Lys
 115 120 125
 Ala Leu Met Ser Leu Ile Arg Gln Asp Val Lys Asn Ser Leu Gln Pro
 130 135 140

Leu Phe Lys His Val Glu Gln Gly Ser Glu Ile Arg Glu Lys Ile Ile
 145 150 155 160
 Cys Phe Leu Arg Asp Lys Val Phe Pro Leu Lys Ala Glu Leu Leu Lys
 165 170 175
 Pro Gln Ala Glu Met Glu Arg Phe Ile Thr Asp Leu Ile Lys Lys Ser
 180 185 190
 Val Gln Asp Val Thr Gly Ser Glu Phe Glu Leu Phe Met Gly Phe Leu
 195 200 205
 Arg Xaa Trp Ser Ile Phe Gly Asp Ser Ala Pro Arg Glu Ser Phe Gln
 210 215 220
 Glu Leu Ile Glu Ile Ile Gln Ala Gln Ala Asp Leu Asn Ser Gln Phe
 225 230 235 240
 Asn Val Ser Asp Ile Asp His Ile Glu Arg Trp Ile Ser Cys Met Tyr
 245 250 255
 Met Ala Leu Pro Ile Phe Met Arg Gly Ala Ser Ala Ser Lys Phe Leu
 260 265 270
 Asn Tyr Phe Val Lys Gln Ile Val Pro Ala Phe Glu Lys Ile Pro Glu
 275 280 285
 Glu Lys Lys Leu Asp Leu Leu Lys Thr Ile Ala Ser Ser Ser Pro Tyr
 290 295 300
 Ala Thr Ala Gln Asp Ser Arg Gln Leu Leu Pro Ser Val Val Gln Leu
 305 310 315 320
 Leu Asn Lys Tyr Met Pro Gly Lys Lys Val Asp Asp Ile Asn His Asn
 325 330 335
 Tyr Val Glu Cys Leu Leu Tyr Thr Tyr His His Leu Ala His Lys Thr
 340 345 350
 Pro Asn Thr Thr Asn Ser Leu Cys Gly Tyr Lys Ile Val Thr Gly Gln
 355 360 365
 Pro Ser Asp Arg Leu Gly Glu Asp Phe Thr Glu His Tyr Lys Asp Phe
 370 375 380
 Thr Glu Arg Leu Thr Gly Thr Glu Glu Thr Val Arg Ala Ala Ser Lys
 385 390 395 400
 Arg Leu Thr Gln Gly Met Ala Asp Phe Ser Lys Ala Ile Ser Ser Ala
 405 410 415
 Lys Thr Glu Glu Glu Lys Thr Lys Ile Lys Gly Asp Gln Gln Thr Ser
 420 425 430
 Thr Arg Thr Met Arg Ser Tyr Asn Asn Ile Leu Ala Met Thr Gln Ser
 435 440 445
 Leu His Ser Lys Ser Pro Leu Phe Ile Gly Asp Lys Lys Ile Thr Leu
 450 455 460
 Ser Trp Met Glu Gln Pro Asn Lys Ala Ala Ala Thr Lys Ala Gly Gly
 465 470 475 480
 Lys Arg Ser Gln Pro Ser Thr Asn Gly Asn Asp Pro Ala Asn Lys Lys
 485 490 495

Gly Arg Gly Gly Met Gln Asn Gln Leu Val Asn Arg Ala Phe Glu Gly
500 505 510

Leu Ser His Val Gly Arg Gly Ser Gly Arg Xaa Arg Gly Xaa Gly Gly
515 520 525

Pro Arg Lys Arg Lys Arg Met Glu Trp Gly Thr Thr Glu Met Thr Trp
530 535 540

Lys Leu Cys Ser Asn Leu Thr Arg Asp Glu Leu Ser Ala Ser Thr Arg
545 550 555 560

Glu

<210> 9
<211> 556
<212> DNA
<213> Oryza sativa

<220>
<221> unsure
<222> (179)

<220>
<221> unsure
<222> (339)

<220>
<221> unsure
<222> (345)

<220>
<221> unsure
<222> (368)

<220>
<221> unsure
<222> (390)

<220>
<221> unsure
<222> (392)

<220>
<221> unsure
<222> (427)

<220>
<221> unsure
<222> (434)

<220>
<221> unsure
<222> (464)

<220>
<221> unsure
<222> (472)

<220>
<221> unsure
<222> (496)

<220>
 <221> unsure
 <222> (518)

<220>
 <221> unsure
 <222> (524)

<220>
 <221> unsure
 <222> (534)

<220>
 <221> unsure
 <222> (551)

<400> 9
 ggcataactaa cccccccaaa tctccacac cgctcccccg ccgccatggc cgcctccgac 60
 gccgacgccg cggaggtcga gcggctctac gagctcggcg agcgctctc ctccgccaaag 120
 gacaagtccc agcacgcggc ggactacgag gcgatcatat cggccgtgaa ggggcaganc 180
 gtgaaggcga agcagctcgc ggcgagctc atcccccgct tcttccggag cttcccggca 240
 ctgccccgc gcgccatgga ggccatgttc gacctcgtcg acatggatga actcgcgact 300
 agaatacaac tattcgtggg ttttcacttc ttgccaaana tgcanaattt gtctcaaaaa 360
 ttgccganat ccttggacaa tccttgcaan tnaggaaaat gtggacgtga tgctgtcata 420
 aagcacngat gtcncttata cggcaggatt taaaattctt gcancctatt angattggat 480
 tcgggatata attctnaaaa ttattgttcc taagaaangc tccngtaaag aaantgtgaa 540
 ctcaacagag ngaaat 556

<210> 10
 <211> 131
 <212> PRT
 <213> Oryza sativa

<220>
 <221> UNSURE
 <222> (45)

<220>
 <221> UNSURE
 <222> (98)

<220>
 <221> UNSURE
 <222> (100)

<220>
 <221> UNSURE
 <222> (108)

<220>
 <221> UNSURE
 <222> (115) .. (116)

<220>
 <221> UNSURE
 <222> (128)

<220>
 <221> UNSURE
 <222> (130)

<400> 10
 Met Ala Ala Ser Asp Ala Asp Ala Ala Glu Val Glu Arg Leu Tyr Glu
 1 5 10 15

Leu Gly Glu Arg Leu Ser Ser Ala Lys Asp Lys Ser Gln His Ala Ala
 20 25 30
 Asp Tyr Glu Ala Ile Ile Ser Ala Val Lys Gly Gln Xaa Val Lys Ala
 35 40 45
 Lys Gln Leu Ala Ala Gln Leu Ile Pro Arg Phe Phe Arg Ser Phe Pro
 50 55 60
 Ala Leu Ala Pro Arg Ala Met Glu Ala Met Phe Asp Leu Val Asp Met
 65 70 75 80
 Asp Glu Leu Ala Thr Arg Ile Gln Leu Phe Val Gly Phe His Phe Leu
 85 90 95
 Pro Xaa Met Xaa Asn Leu Ser Gln Lys Leu Pro Xaa Ser Leu Asp Asn
 100 105 110
 Pro Cys Xaa Xaa Gly Lys Cys Gly Arg Asp Ala Val Ile Lys His Xaa
 115 120 125
 Cys Xaa Leu
 130

<210> 11
 <211> 749
 <212> DNA
 <213> Glycine max

<220>
 <221> unsure
 <222> (678)

<220>
 <221> unsure
 <222> (724)

<220>
 <221> unsure
 <222> (745)

<400> 11
 gcgctggcta ctgtttatag tttactgtaa actgtgttta cgttgtgtcg tgtggcgtgt 60
 tcagttaggt aagggaact cgtccccatc gaagagctta cttgacctcg caccacggaa 120
 tcgttccta ctaattcaac tcaacaacac tatcgtctcc attcactagt tagaaacgtg 180
 cgttccaatg tctgatcctg ccgaagaggc tgctttcatc gagaagctct acgaatacgg 240
 cgagcaactc aacaacacta tcgtctccat tcactagtta gaaacgtgcg ttccaatgtc 300
 tgatcctgcc gaagaggctg ctttcatcga gaagctctac gaatacggcg agcaactcaa 360
 caatgccaaag gacaagtcgc agaattgtgca ggattaccag ggaatcatag atgcggcgaa 420
 gacgagtgtg aaggcgaagc agctcgctgc acagctgatt cccaggttct acaagttctt 480
 tcctgacctt tctagccctg ctctcgatgc acatcttgat ttggttgagg ctgaagaact 540
 cgggggttcga gtgcaagcaa ttagaggtct gcctcttttt tgtaaggata cacctgagaa 600
 tattgggaag atggttgata ttcttgatgca aattcttggg tctgaggaat ttgtggagcg 660
 tgatgcagta cataaggntc ttaagtcctt tgctgaggca aggatgtcaa aagcttcctt 720
 gacngctttg ttaagcacia ttgnaagg 749

<210> 12
 <211> 131
 <212> PRT
 <213> Glycine max

<220>
 <221> UNSURE
 <222> (128)

<400> 12
 Met Ser Asp Pro Ala Glu Glu Ala Ala Phe Ile Glu Lys Leu Tyr Glu
 1 5 10 15
 Tyr Gly Glu Gln Leu Asn Asn Ala Lys Asp Lys Ser Gln Asn Val Gln
 20 25 30
 Asp Tyr Gln Gly Ile Ile Asp Ala Ala Lys Thr Ser Val Lys Ala Lys
 35 40 45
 Gln Leu Ala Ala Gln Leu Ile Pro Arg Phe Tyr Lys Phe Phe Pro Asp
 50 55 60
 Leu Ser Ser Pro Ala Leu Asp Ala His Leu Asp Leu Val Glu Ala Glu
 65 70 75 80
 Glu Leu Gly Val Arg Val Gln Ala Ile Arg Gly Leu Pro Leu Phe Cys
 85 90 95
 Lys Asp Thr Pro Glu Asn Ile Gly Lys Met Val Asp Ile Leu Val Gln
 100 105 110
 Ile Leu Gly Ser Glu Glu Phe Val Glu Arg Asp Ala Val His Lys Xaa
 115 120 125
 Leu Lys Ser
 130

<210> 13
 <211> 115
 <212> PRT
 <213> Arabidopsis thaliana

<400> 13
 Met Val Lys Ser Thr Ser Lys Asp Ala Gln Asp Leu Phe His Ser Leu
 1 5 10 15
 His Ser Ala Tyr Thr Ala Thr Pro Thr Asn Leu Lys Ile Ile Asp Leu
 20 25 30
 Tyr Val Cys Phe Ala Val Phe Thr Ala Leu Ile Gln Val Ala Tyr Met
 35 40 45
 Ala Leu Val Gly Ser Phe Pro Phe Asn Ser Phe Leu Ser Gly Val Leu
 50 55 60
 Ser Cys Ile Gly Thr Ala Val Leu Ala Val Cys Leu Arg Ile Gln Val
 65 70 75 80
 Asn Lys Glu Asn Lys Glu Phe Lys Asp Leu Ala Pro Glu Arg Ala Phe
 85 90 95
 Ala Asp Phe Val Leu Cys Asn Leu Val Leu His Leu Val Ile Ile Asn
 100 105 110
 Phe Leu Gly
 115

<210> 14
 <211> 114
 <212> PRT
 <213> Oryza sativa

<400> 14

Met Pro Arg Ala Thr Ser Asp Ala Lys Leu Leu Ile Gln Ser Leu Gly
 1 5 10 15
 Lys Ala Tyr Ala Ala Thr Pro Thr Asn Leu Lys Ile Ile Asp Leu Tyr
 20 25 30
 Val Val Phe Ala Val Ala Thr Ala Leu Ile Gln Val Val Tyr Met Gly
 35 40 45
 Ile Val Gly Ser Phe Pro Phe Asn Ser Phe Leu Ser Gly Val Leu Ser
 50 55 60
 Cys Ile Gly Thr Ala Val Leu Ala Val Cys Leu Arg Ile Gln Val Asn
 65 70 75 80
 Lys Asp Asn Lys Glu Phe Lys Asp Leu Pro Pro Glu Arg Ala Phe Ala
 85 90 95
 Asp Phe Val Leu Cys Asn Leu Val Leu His Leu Val Ile Met Asn Phe
 100 105 110

Leu Gly

<210> 15

<211> 117

<212> PRT

<213> Pisum sativum

<400> 15

Met Ala Lys Thr Ser Ser Thr Thr Lys Asp Ala Gln Asp Leu Phe His
 1 5 10 15
 Ala Ile Trp Ser Ala Tyr Ser Ala Thr Pro Thr Asn Leu Lys Ile Ile
 20 25 30
 Asp Leu Tyr Val Val Phe Ala Val Phe Thr Ala Leu Leu Gln Asp Val
 35 40 45
 Tyr Met Ala Leu Val Gly Pro Phe Pro Phe Asn Ser Phe Leu Ser Gly
 50 55 60
 Val Leu Ser Cys Val Gly Thr Ala Val Leu Ala Val Cys Leu Arg Ile
 65 70 75 80
 Gln Val Asn Lys Glu Asn Lys Glu Phe Lys Asp Leu Gly Pro Glu Arg
 85 90 95
 Ala Phe Ala Asp Phe Val Leu Cys Asn Leu Val Leu His Leu Val Ile
 100 105 110

Met Asn Phe Leu Gly
115

<210> 16

<211> 528

<212> PRT

<213> Homo sapiens

<400> 16

Met Gly Leu Ser Leu Thr Met Pro Thr Val Glu Glu Leu Tyr Arg Asn
 1 5 10 15

Tyr Gly Ile Leu Ala Asp Ala Thr Glu Gln Val Gly Gln His Lys Asp
 20 25 30
 Ala Tyr Gln Val Ile Leu Asp Gly Val Lys Gly Gly Thr Lys Glu Lys
 35 40 45
 Arg Leu Ala Ala Gln Phe Ile Pro Lys Phe Phe Lys His Phe Pro Glu
 50 55 60
 Leu Ala Asp Ser Ala Ile Asn Ala Gln Leu Asp Leu Cys Glu Asp Glu
 65 70 75 80
 Asp Val Ser Ile Arg Arg Gln Ala Ile Lys Glu Leu Pro Gln Phe Ala
 85 90 95
 Thr Gly Glu Asn Leu Pro Arg Val Ala Asp Ile Leu Thr Gln Leu Leu
 100 105 110
 Gln Thr Asp Asp Ser Ala Glu Phe Asn Leu Val Asn Asn Ala Leu Leu
 115 120 125
 Ser Ile Phe Lys Met Asp Ala Lys Gly Thr Leu Gly Gly Leu Phe Ser
 130 135 140
 Gln Ile Leu Gln Gly Glu Asp Ile Val Arg Glu Arg Ala Ile Lys Phe
 145 150 155 160
 Leu Ser Thr Lys Leu Lys Thr Leu Pro Asp Glu Val Leu Thr Lys Glu
 165 170 175
 Val Glu Glu Leu Ile Leu Thr Glu Ser Lys Lys Val Leu Glu Asp Val
 180 185 190
 Thr Gly Glu Glu Phe Val Leu Phe Met Lys Ile Leu Ser Gly Leu Lys
 195 200 205
 Ser Leu Gln Thr Val Ser Gly Arg Gln Gln Leu Val Glu Leu Val Ala
 210 215 220
 Glu Gln Ala Asp Leu Glu Gln Thr Phe Asn Pro Ser Asp Pro Asp Cys
 225 230 235 240
 Val Asp Arg Leu Leu Gln Cys Thr Arg Gln Ala Val Pro Leu Phe Ser
 245 250 255
 Lys Asn Val His Ser Thr Arg Phe Val Thr Tyr Phe Cys Glu Gln Val
 260 265 270
 Leu Pro Asn Leu Gly Thr Leu Thr Thr Pro Val Glu Gly Leu Asp Ile
 275 280 285
 Gln Leu Glu Val Leu Lys Leu Leu Ala Glu Met Ser Ser Phe Cys Gly
 290 295 300
 Asp Met Glu Lys Leu Glu Thr Asn Leu Arg Lys Leu Phe Asp Lys Leu
 305 310 315 320
 Leu Glu Tyr Met Pro Leu Pro Pro Glu Glu Ala Glu Asn Gly Glu Asn
 325 330 335
 Ala Gly Asn Glu Glu Pro Lys Leu Gln Phe Ser Tyr Val Glu Cys Leu
 340 345 350
 Leu Tyr Ser Phe His Gln Leu Gly Arg Lys Leu Pro Asp Phe Leu Thr
 355 360 365

Ala Lys Leu Asn Ala Glu Lys Leu His Glu Ser Lys Ile Arg Leu Gln
 370 375 380
 Tyr Phe Ala Arg Gly Leu Gln Val Tyr Ile Arg Gln Leu Arg Leu Ala
 385 390 395 400
 Leu Gln Gly Lys Thr Gly Glu Ala Leu Lys Thr Glu Glu Asn Lys Ile
 405 410 415
 Lys Val Val Ala Leu Lys Ile Thr Asn Asn Ile Asn Val Leu Ile Lys
 420 425 430
 Asp Leu Phe His Ile Pro Pro Ser Tyr Lys Ser Thr Val Thr Leu Ser
 435 440 445
 Trp Lys Pro Val Gln Lys Val Glu Ile Gly Gln Lys Arg Ala Ser Glu
 450 455 460
 Asp Thr Thr Ser Gly Ser Pro Pro Lys Lys Ser Ser Ala Gly Pro Lys
 465 470 475 480
 Arg Met Pro Gly Arg Phe Ile Thr Leu Pro Val Gly Asn Ile Ala Ala
 485 490 495
 Ile Trp Ala Thr Leu Ile Met Arg Gly Ala Phe Arg Gly Ser Lys Trp
 500 505 510
 Pro Arg Leu Gly His Thr Lys Glu Ile Val Val Gly Glu Arg Leu Tyr
 515 520 525

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/82, 15/29, 5/10, C07K 14/415, C12Q 1/68		A3	(11) International Publication Number: WO 99/47689 (43) International Publication Date: 23 September 1999 (23.09.99)
(21) International Application Number: PCT/US99/05227 (22) International Filing Date: 10 March 1999 (10.03.99) (30) Priority Data: 60/078,144 16 March 1998 (16.03.98) US (71) Applicant (for all designated States except US): E.I. DU PONT DE NEMOURS AND COMPANY [US/US]; 1007 Market Street, Wilmington, DE 19898 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): CAHOON, Rebecca, E. [US/US]; 2331 West 18th Street, Wilmington, DE 19806 (US). CAIMI, Perry, G. [US/US]; 7 Holly Drive, Kennett Square, PA 19348 (US). KLEIN, Theodore, M. [US/US]; 2229 Rosewood Drive, Wilmington, DE 19810 (US). MORGANTE, Michele [IT/US]; 2306 Lorelei Lane, Wilmington, DE 19810 (US). SAKAI, Hajime [DE/US]; 105 Banbury Drive, Wilmington, DE 19803 (US). SHEN, Jennie, Bih-Jien [US/US]; 15 Bromley Court, Wilmington, DE 19810 (US). TINGEY, Scott, V. [US/US]; 35 Somerset Court, Elkton, MD 21921 (US).			(74) Agent: MAJARIAN, William, R.; E.I. du Pont de Nemours and Company, Legal Patent Records Center, 1007 Market Street, Wilmington, DE 19898 (US). (81) Designated States: AL, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GD, GE, HR, HU, ID, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> (88) Date of publication of the international search report: 25 November 1999 (25.11.99)
(54) Title: INHIBITORS OF APOPTOSIS PROTEINS IN PLANTS			
(57) Abstract <p>This invention relates to an isolated nucleic acid fragment encoding an apoptosis inhibitory protein. The invention also relates to the construction of a chimeric gene encoding all or a portion of the apoptosis inhibitory protein, in sense or antisense orientation, wherein expression of the chimeric gene results in production of altered levels of the apoptosis inhibitory protein in a transformed host cell.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

International Application No
PC 1/US 99/05227

A. CLASSIFICATION OF SUBJECT MATTER IPC 6 C12N15/82 C12N15/29 C12N5/10 C07K14/415 C12Q1/68		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 6 C12N C07K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	TANAKA, Y., ET AL.: "dad-1, a putative programmed cell death suppressor gene in rice" PLANT CELL PHYSIOL., vol. 38, no. 3, 1997, pages 379-383, XP002085156 abstract, page 379, right column; page 383; Fig. 1 + 2	1-3, 9-11, 23-27
X	--- GALLOIS, P., ET AL.: "an Arabidopsis thaliana cDNA complementing a hamster apoptosis suppressor mutant" THE PLANT JOURNAL, vol. 11, no. 6, 1997, pages 1325-1331, XP002106087 abstract; page 1326; Fig. 1+2 --- <div style="text-align: center;">-/--</div>	1-3, 9-11,23
<div style="display: flex; justify-content: space-between;"> <input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input type="checkbox"/> Patent family members are listed in annex. </div>		
* Special categories of cited documents :		
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>*A* document defining the general state of the art which is not considered to be of particular relevance</p> <p>*E* earlier document but published on or after the international filing date</p> <p>*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>*O* document referring to an oral disclosure, use, exhibition or other means</p> <p>*P* document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>*Z* document member of the same patent family</p> </div> </div>		
Date of the actual completion of the international search <div style="text-align: center; font-weight: bold;">5 July 1999</div>	Date of mailing of the international search report <div style="text-align: center; font-weight: bold;">07.10.99</div>	
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016	Authorized officer <div style="text-align: center; font-weight: bold;">Holtorf, S</div>	

Form PCT/ISA/210 (second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

International Application No

PC 1/US 99/05227

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>ORZAEZ, D., ET AL.: "the plant homologue of the defender against apoptotic death gene is down-regulated during senescence of flower petals"</p> <p>FEBS LETTERS, vol. 404, 1997, pages 275-278, XP002106088 abstract; page 275; Fig. 2; page 276, right column; page 278</p>	1-3, 9-11,23, 25,27
A	<p>APTE, S.S., ET AL.: "the highly conserved defender against the death 1 (DAD1) gene maps to human chromosome 14q11-q12 and mouse chromosome 14 and has plant and nematode homologs"</p> <p>FEBS LETTERS, vol. 363, 1995, pages 304-306, XP002106089 abstract; fig. 1</p>	1-3,6, 9-11, 23-27
A	<p>SUGIMOTO, A., ET AL.: "dad-1, an endogenous programmed cell death suppressor in Caenorhabditis elegans and vertebrates"</p> <p>THE EMBO JOURNAL, vol. 14, no. 18, 1995, pages 4434-4441, XP002106090 figure 3</p>	1-3,6, 9-11, 23-27
A	<p>KELLEHER, D., ET AL.: "DAD1, the defender against apoptotic cell death, is a subunit of the mammalian oligosaccharyltransferase"</p> <p>PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE USA, vol. 94, May 1997 (1997-05), pages 4994-4999, XP002106091 abstract; pages 4994, right column; page 4998, right column; page 4999</p>	1-3,6, 9-11, 23-27
A	<p>MASTRANGELO A J ET AL: "Overcoming apoptosis: new methods for improving protein-expression systems"</p> <p>TRENDS IN BIOTECHNOLOGY, vol. 16, no. 2, 1 February 1998 (1998-02-01), page 88-95 XP004107048 page 93, right-hand column</p>	1-3,6, 9-11, 23-27

INTERNATIONAL SEARCH REPORT

national application No.

PCT/US 99/ 05227

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

3,6 completely; 1,2,9-11,23-27 partially

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

1. Claims: claims 3,6 completely; 1,2,9-11,23-27 partially

Isolation and characterization of a Phaseolus-specific DAD1-related cDNA and corresponding amino acid sequence and recombinant expression of the same.

2. Claims: claims 4,7 completely; 1,2,9-11,23-27 partially

Isolation and characterization of a Maize-specific DAD1-related cDNA and corresponding amino acid sequence and recombinant expression of the same.

3. Claims: claims 5,8 completely; 1,2,9-11,23-27 partially

Isolation and characterization of a Soybean-specific DAD1-related cDNA and corresponding amino acid sequence and recombinant expression of the same.

4. Claims: claims 12-22 completely; 23-27 partially

Isolation and characterization of plant-specific ACC-11 related cDNAs and corresponding amino acid sequences and recombinant expression of the same.